BERICHTE
aus dem Fachbereich Geowissenschaften
der Universität Bremen

No. 260

Westphal, H., A. Freiwald, T. Hanebuth, M. Eisele, K. Gürs,
K. Heindel, J. Michel. J. V. Reumont

# REPORT AND PRELIMINARY RESULTS OF POSEIDON CRUISE 346, MACUMA

**- Modern atypical tropical carbonates in the upwelling off Mauretania -,**

**LAS PALMAS (SPAIN) - LAS PALMAS (SPAIN),
28.12.2006 - 15.1.2007.**

Berichte, Fachbereich Geowissenschaften, Universität Bremen, No. 260, 49 pages,
Bremen 2007

ISSN 0931-0800

The "Berichte aus dem Fachbereich Geowissenschaften" are produced at irregular intervals by the Department of Geosciences, Bremen University.

They serve for the publication of experimental works, Ph.D.-theses and scientific contributions made by members of the department.

Reports can be ordered from:

Monika Bachur

Forschungszentrum Ozeanränder, RCOM

Universität Bremen

Postfach 330 440

**D 28334 BREMEN**

Phone: (49) 421 218-65516

Fax: (49) 421 218-65515

e-mail: MBachur@uni-bremen.de

Citation:
Westphal, H. and cruise participants
Report and preliminary results of Poseidon cruise 346, MACUMA. Las Palmas (Spain) – Las Palmas (Spain), 28.12.2006 – 15.1.2007. 49 pages, Bremen 2007.
Berichte, Fachbereich Geowissenschaften, Universität Bremen, No. 260, 49 pages. Bremen 2007.

**ISSN 0931-0800**

# Cruise Report

# Poseidon 346

# MACUMA

## Modern atypical tropical carbonates in the upwelling off Mauritania

Integrating carbonates, siliciclastics and deep-water reefs for understanding a complex environment




Las Palmas – Las Palmas
28.12.2006-15.1.2007

Edited by Hildegard Westphal
With contributions of André Freiwald, Till Hanebuth, Markus Eisele, Karl Gürs, Katrin Heindel, Julien Michel, Jonas v Reumont

## CONTENTS

**1. Abstract** **3**
**2. Participants** **4**
**3. Research Program** **5**
**4. Cruise Narrative** **8**
**5. Scientific Equipment** **12**
**6. Preliminary Results** **16**
6.1. Carbonates of the Gulf d'Arguin 16
6.2. Mud belt of Banc d'Arguin 22
6.3. Coral mounds 27
6.4. CTD measurements 30
**7. Weather and Meteorological Conditions** **35**
**8. Acknowledgements** **36**
**9. References** **36**

**Appendices**

**A1. Weekly reports**
**A2. Technical report**
**A3. Stations book**

# 1. Abstract

Poseidon cruise 346 was a multidisciplinary expedition (sedimentology, biology, oceanography) aiming at better understanding the complex sedimentary system of the upwelling zone off Mauritania. The shallow Banc d'Arguin in the North of Mauritania and the continental slope West of Nouakchott were studied in order to address three main, interrelated objectives:

1. The high-nutrient tropical carbonates of the Gulf d'Arguin
2. The siliciclastic sedimentary record of the Banc d'Arguin and Mauritanian outer shelf
3. The deep-water reefs along the continental slope.

The sedimentary system of Banc d'Arguin is one of the rare examples in the modern world where high-nutrient yet warm waters host carbonate production. The siliciclastic record with sedimentation rates of up to 1 m per 1 ka allows for high-resolution paleoclimate reconstruction of the Late Holocene. The deep-water reefs are the southernmost known occurrence of such ecosystems in the Eastern North Atlantic. The aim is to decipher this complex sedimentary system and its climatic and oceanographic record.

Our work is integrated in a larger study on high-nutrient tropical carbonates (DFG; HW), Northwest African mud belts (DFG; TH), and comprehensive EU- and DFG-funded deep-water reef work (AF).

# 2. Participants

| Name | Discipline | Institution |
|---|---|---|
| Eisele, Markus | Sedimentology | GeoB / RCOM |
| Freiwald, André | Paleontology | IPAL |
| Gürs, Karl | Paleontology | LANUSH |
| Hanebuth, Till | Sedimentology | GeoB / RCOM |
| Heindel, Katrin | Sedimentology | GeoB |
| Lackmann, Oliver | Technical Support | GeoB / RCOM |
| Max, Lars | Sedimentology | GeoB / RCOM |
| Peichert, Nadine | Sedimentology | GeoB |
| v. Reumont, Jonas | Paleoceanography | IFM-Geomar |
| Westphal, Hildegard* | *Chief Scientist* | GeoB |
| Abed, Jemal Ould | Observer | IMROP |

| | |
|---|---|
| GeoB | Department of Geosciences, Bremen University |
| RCOM | Research Center Ocean Margins, Bremen University |
| LANUSH | Landesamt für Natur und Umwelt, Schleswig-Holstein |
| IFM-Geomar | Leibniz-Institut für Meeresforschung und Zentrum für Marine Geowissenschaften, Kiel University |
| IMROP | Institut Mauritanien de Recherches Océanographiques et des Pêches, Nouadhibou, Mauritania |

Fig. 1: Scientific team of RV Poseidon cruise 346 (JvR, OL, KH, ME, TH, HW, ML, KG, AF; NP not on picture).

*) FB 5, Geowissenschaften, MARUM-Gebäude, Leobener Straße, 28359 Bremen, Deutschland, hildegard.westphal@uni-bremen.de

# 3. Research Program

## Research Objectives

The MACUMA research cruise dealt with the mixed carbonate-siliciclastic sedimentary system on the shelf and shelf slope off Mauretania (Banc d'Arguin and west of Nouakchott) (see map with cruise track, Fig. 1). MACUMA stands for Modern Atypical tropical Carbonates in the influence of Upwelling off MAuritania. With a latitudinal extension from 17° to 21° N, the study area is located in the tropical climate belt. It is however under the influence of the southward-directed Canary Current and an oceanic upwelling front, modifying the conditions in complex ways. The sedimentary system reflects this modification in that it lacks typical tropical sediment-forming elements such as zooxanthellate coral reefs or calcareous green algae.

The research objectives for the research cruise were:

1. Study of the modern atypical tropical carbonate sedimentary system of the Gulf d'Arguin;
2. Study of the deep-water coral reefs on the continental slope;
3. Study of the Quaternary climate evolution as recorded in the fluvial and eolian influence on the siliciclastic and carbonate sedimentary system of the Gulf d'Arguin.




Fig. 2: Map of study area and cruise track.

(1) The carbonate sedimentary system of the Banc d'Arguin is a rare example of such atypical tropical carbonates. The study area was chosen, because in contrast to most other upwelling areas in the modern world, water temperatures on the shallow shelf are tropical and do not drop below the critical benchmark of 18°C. Nevertheless the waters are highly enriched in nutrients and are very productive. This combination of warm, nutrient rich waters, even though being uncommon in the modern world, might have been more widespread in the past. For reliable interpretation of ancient occurrences of such carbonate sediments, modern analogues need to be comprehensively studied. In addition, warm, nutrient-rich waters might become more widespread in the future as a result of global warming, thoroughly changing the carbonate producing systems and thus the physical behavior and morphology of shallow-water areas. One aim of the cruise was therefore to understand the complex reaction of the ecosystem and the resulting sedimentary system to such elevated-nutrient, warm waters.

(2) At the shelf slope, largely unknown deep-water coral reefs represent another peculiarity of the sedimentary system. These deep-water reefs compose the southernmost part of the irregular reef belt from northern Norway southward along the eastern Atlantic shelf slope. These deep-water reefs represent important fish breeding environments and therefore are not only scientifically but also commercially of high interest. One objective of the cruise was to study the faunal composition of these southernmost reef mounds and to compare them with the North-South trends along the North Atlantic deep-water reef belt.

(3) The mixed carbonate-siliciclastic system of the middle and southern Banc d'Arguin composes a unique high-resolution climate archive of the Quaternary (Hanebuth, in prep.). It records the interplay between eolian (dust) and fluvial influx of siliciclastic sediment through time. The aims of the cruise therefore included to answer the question whether the sedimentary record is continuous and allows for reconstructing Late Holocene climate evolution with a high resolution, and to reconstruct paleo-river activity in the study area that might have drained the hinterland into the Golf d'Arguin during the Holocene Climate Optimum.

The study area spans from the Baie du Lévrier east of Cap Blanc along the shelf bordering the Banc d'Arguin and further south to the deep-water reef mounds off Nouakchott. For all three objectives, examination of sedimentology, taxonomy, geochemistry, and oceanography is currently being undertaken. All three objectives will be studied as part of a comprehensive approach that will integrate the three aspects of the sedimentary system.

## Scientific Program and Sampling Strategy

The scientific program was designed to provide

- Surface sediment sample coverage (box corer, grab sampler) of the shallow Baie du Levrier where possible with RV Poseidon;
- Surface sediment sample coverage (box corer, grab sampler) of the 20-30 m isobath along the western slope of the Banc d'Arguin;
- A small number of vibro-cores for these areas in order to extend the record of carbonate production back in time (vibro-corer needed for penetrating the coarse shell layers);

- CTD and water sampling in E-W transects off Banc d'Arguin for a better understanding of the influence of the different water masses on carbonate production on the banc;
- Gravity cores and vibro-cores of the mud wedges off the southern Banc d'Arguin (80-100 m water depth) in order to complement the existing data set (Hanebuth, subm.) for climate reconstruction of the late Holocene;
- Gravity cores and surface samples of the deep-water reef mounds west of Nouakchott for a first faunistic and sedimentary study of these southern deep-water reefs of the North-Atlantic deep-water reef chain;
- CTD and water sampling west and on the deep-water reef mounds to gain first information on the oceanographic conditions related to these mounds.

# 4. Cruise Narrative

H. Westphal

The RV Poseidon left the harbor of Las Palmas on the 28th at 9:00 local time and started the transit to the study area off Mauritania. The team assembled geologists and paleontologists from University of Bremen, University of Erlangen, and the Landesamt für Natur und Umwelt (Schleswig-Holstein). Additionally, a guest scientist from IMROP (Nouadhibou, Mauritania) was on board the vessel.

On the 30th we had arrived at the study area and at 22:45 the Poseidon was on the first station where a CTD cast was carried out in 1550 m water depth west of the Banc d'Arguin. The temperature and water distribution of the water column clearly reflected the Canary Current. Work on the shelf off the Banc d'Arguin was commenced on the 31st, when at five stations box cores and gravity cores were taken. This southern part of the shelf is characterized by a fine-grained mud wedge (Timiris Mud Wedge, TMW) in roughly 80 m water depth that is composed of dark sediment with coarse carbonate shells. During the night from Dec. 31st to Jan. 1st an echo sounder section was measured for reconnaissance of the shelf morphology that is disrupted by submarine canyons.

Coring work on the 1st of January continued ocean-ward of the TMB. The sediment recovered by box coring and gravity coring again was very fine-grained and dark and had a pronounced $H_2S$ smell. In a water depth of 370 m we found fragments of the deep-water coral *Madrepora* as well as solitary corals. Later that day we started a grab sampler profile northward along the 20 m isobath of the Banc d'Arguin that should add up to some 20 samples along the entire N-S extension of the Banc during the cruise. Even in this shallower water depth, in the northern part of this profile the sediment was dark colored, however well-sorted quartz sand was identified as major component in some samples, pointing to possible submerged eolian dunes. The sediment was rich in worm tubes and live worms and $H_2S$ smell. The high productivity of the area was also manifested in considerable concentrations of otoliths in the sediment and also in the fact that several box cores brought live fish on deck.

During the night to the 2nd of January, wind strength increased considerably to up to Wind Force 8 (Harmattan; see below). Therefore the planned CTD stations were cancelled. In an attempt to mitigate the effects of the storm, we moved closer to the coastline near Cap Blanc in the North of the Banc d'Arguin. Shortly after noon the winds ceased and grab sampling and box coring of the shallow waters of the northern Banc revealed clean, lose, and coarse carbonate sands and shell deposits. Two attempts to core this coarse carbonate sediment with the vibro-corer were unsuccessful due to technical problems. During the night, five CTD casts were measured in a section from the northern Banc to the open ocean. The same stations were sampled by means of box corer and gravity corer the next day (Jan. 3rd), revealing dark fine-grained sediment. On the surface of the box cores were abundant live ophiurids and crinoids. In the night, the 20 m isobath profile was continued up to off Cap Blanc with 10 grab sample stations. The sediment showed a general trend to cleaner, carbonate-rich, coarser sediment northward and

finer, dark sediment southward. For one station off Cap Blanc, the sediment was dominated by well-sorted quartz sand reminiscent of desert sand.

On the 4th of Jan. we continued our work on the southern Banc d'Arguin in a presumed paleo-valley setting (Arguin Mud Wedge; AMW). Due to rough conditions, the vibro-corer could only be used after noon; during the morning, box coring and grab sampling was undertaken. During the night, CTD casts were measured, and on the 5th of Jan, several grab samples were taken along the 20 m isobath profile to increase data density and to wait for calm conditions in order to attempt additional vibro-coring in the paleo-valley setting. Three cores were successfully recovered during the afternoon.

In the evening of the 5th of Jan., we left the Banc d'Arguin and headed south for deep-water coral targets South of Nouakchott. During the transit, several CTD casts were measured. The Poseidon arrived in the coral mound province (Banda Mounds) in the morning of Jan. 6th. In sight of the Chinguetti Oil Field we spent the entire day with reconnaissance by means of echo sound in order to exactly locate the coral mounds and gain information on their morphology. Reflections typical for coral mounds were observed with steep slopes and soft reflectors sitting on sharp basal reflectors. The echo sounder profiling, however, was strongly disturbed by large herds of dolphins. During the following night, CTD casts were measured along the Banda Mound chain. On Jan 7th, a series of box cores and gravity cores was recovered from top mound and slope locations. Even though the samples were very rich in deep-water corals, only one live *Lophelia* was recovered. During the night, grab sampling was undertaken to gain samples from a transect towards the coastline. Box coring and gravity coring of coral mounds was continued on Jan. 8th again recovering abundant deep-water corals (*Lophelia*, *Madrepora*, and *Dendrophyllia*), and the deep-water reef bivalve *Acesta*.

During the night, the Poseidon proceeded to the northern Timiris Mound province. Echo sounder profiles were carried out and a series of CTD casts were measured. On the 9th of Jan. in the morning, box cores and gravity cores of the Timiris Mounds were recovered. Then the vessel left the coral mound provinces and returned to the southern Banc d'Arguin, measuring two CTD profiles along the way. In the afternoon, East of the TMB, a dinghy was launched in order to gain sample material from the shallow Banc d'Arguin. The bathymetry in this southern part of the Banc is rather abrupt, and the dinghy only went for some 1.5 nm before reaching a water depth of roughly 3.5 m. Currents were strong, and due to the extremely calm sea, morphological edges of the shallow sea floor were clearly visible. Sediment was dredged with a simple bucket, revealing the first occurrences of red algae encountered during this cruise. During the night, a CTD cast was measured seaward of the dinghy station, and a series of grab samples were taken East of the TMB as the southernmost extension of the 20 m isobath profile along the edge of Banc d'Arguin.

In the morning of the 10th of Jan., the Baie du Levrier was reached, the northernmost part of Banc d'Arguin, sheltered East of Cap Blanc. The sea was calm and it was slightly raining. A series of grab samples was taken in 10-20 m water depth. The sediment was clean carbonate sand dominated by bivalves. Two attempts were made to recover vibro-cores. The first vibro-corer station was located in the entrance of Baie du Levrier in 12 m water depth. Due to the strong tidal currents, the station was planned for stillwater at 10:00. The corer recovered 5 m of coarse

carbonate shell sediments. At the other station slightly Southwest of the tip of Cap Blanc, the vibro-corer hit a hard surface and was damaged.

During the night, CTD stations were measured along a section in westward direction off Cap Blanc down to water depths of 498 m. In the morning of the 11th of Jan., the weather was humid and rainy. The shallower of the CTD stations of the night were targeted for sediment sampling; grab sampling and box coring took place while the vibro-corer repair was completed. A vibro-core with well-sorted fine sand, shell rich sediment was recovered. As the sea got rougher, the next vibro-corer stations were abandoned, and grab sampling and box coring were attempted. After the weather continued to deteriorate and reached Wind Force 9, at 11:00 all work had to stop. The weather forecast predicted continuing poor conditions, therefore the Poseidon started her transit back to the port in Las Palmas where it arrived on the 15th of Jan. at 9:00 local time.




Fig. 3: Map of the stations during cruise POS 346. light grey: CTD and water samples; black: grab samples and box cores only; dark grey: gravity cores or vibro-cores (plus grab samples and/or box cores).

# 5. Scientific Equipment

M. Eisele, K. Gürs, T. Hanebuth, K. Heindel, J. v. Reumont, H. Westphal

## List of total number of operations

| Equipment | Number of stations/sections/profiles | Successful runs |
|---|---|---|
| Echo sounder sections | 4 | 4 |
| CTD casts / rosette | 34 | 32 |
| Van Veen grab | 62 | 59 |
| Box corer | 35 | 31 |
| Gravity corer | 19 | 19 |
| Vibro-corer | 12 | 8 |

Table 1: Number of stations per tool; detailed list of all stations see appendix.

## Giant Box-Corer

The Giant Box-Corer (50 cm x 50 cm x 50 cm) was a main sampling tool for surface sediments (Fig. 4).

Fig. 4: Giant Box-Corer used during POS 346. Left: Box corer with core is brought back on deck. Right: Sub-sampling of the box core with core liners.

The box corer was run at a 35 stations, a total of 34 box cores were recovered. The recovery was strongly dependent on the sediment (grain size), whereby fine-grained sediment resulted in cores of 40-50 cm length. Box cores taken in coarse shell sediments typical for the shallow water settings of Baie du Levrier only recovered 10-30 cm. The coarse layers hindered penetration of the box corer into the sediment. Sediments from sites situated in deeper water settings and in the southern part of the Banc d'Arguin were dominated by silt-sized grains and showed a recovery of 40-50 cm.

The on-board protocol for the box cores included description (color, grain sizes, fauna, sediment structures) and sampling of the surface. The live macro-fauna was collected and preserved in ethanol (30 %) for taxonomic studies. Two 200 $cm^2$ sub-samples were taken for micropaleontological studies. One of these was stained with a solution of rose Bengal (1 g in 1 l of ethanol) in order to allow distinction of live micro-fauna. Two cores were taken from each box-core. The vertical section was then described, and bulk samples were taken from each visible layer. The remaining sediment was wet-sieved layer by layer with mesh-sizes of 1 cm and 2 cm. The sieved sediment fractions were dried. Digital photography of the surface, the vertical section, and the coarse fraction were taken for each box core.

## Van Veen Grab

The grab sampler (Fig. 5) was used for exploratory sampling prior to employing the various corers, and for surface sediments in comparably high resolution, and as reconnaissance tool for gaining information on sediment types before operating box corer, gravity corer, or vibro-corer. For high-resolution surface sampling, the grab sampler was mainly used during the nights. A total of 62 grab sampling stations were run, 59 of which recovered surface sediment. The grab sample was documented photographically and then sub-sampled (three bulk samples, two syringe samples, live fauna in ethanol). The remaining bulk sediment was sieved with mesh widths of 1 cm and 2 cm.

Fig. 5: Van Veen Grab used during POS 346.

## Gravity Corer

The Gravity Corer was used where the van Veen grab indicated soft and fine sediment conditions, that is, at sites in the South of Banc d'Arguin and at deeper sites. A 6 m core barrel 6 m was used with a weight of 1.800 kg. A total of 19 gravity cores stations were run, all of which recovered cores. An average of 466 cm core length (215-586 cm) was recovered. The cores were cut in one-meter segments and were opened in the laboratory in Bremen. The gravity cores from Banc d'Arguin are stored at Bremen University, the gravity cores from the deep-water coral reef stations are stored at Erlangen University.

## Vibro-corer

At stations judged unsuitable for using the gravity corer (danger of damaging the core barrel), the vibro-corer was used (VKG-5 produced by Th. Schmidt, Rostock, Germany; Fig. 6). It proved successful for recovering shelf sediment successions consisting either of homogeneous coarse-grained sediment, or of fine-grained sediment with interbedded coarse layers. Expected coarser sediments included paleo-coastal deposits, thick shell horizons, and paleosols that cannot be penetrated by gravity corers but can be successfully cored by vibro-systems.

The vibro-corer provided a maximum core length of 500 cm and had a diameter of 10 cm. With the available 175-m long electrical cable, which is run parallel to the steel wire by hand, coring in a maximum water depth of 140 m was possible during calm weather and wave conditions.

At a total of 12 stations the vibro-corer was run, and cores were recovered at eight stations. Technical problems and recurring rough conditions prevented a more frequent usage, however where successfully run, the vibro-corer had extremely high and complete recovery. An average of 454 cm of sediment was recovered (284-491 cm). Once the cores were on deck, the plastic liner was cut into 1 m segments, closed with caps and labeled according the scheme applied for all GeoB cores that are stored at the University of Bremen (MARUM).

Fig. 6: Vibro-corer VKG-5 on the way down.

## CTD and Rosette

The deployed CTD system was a Sea-Bird “SBE 9plus” underwater unit in combination with a Sea-Bird “SBE 11plus” deck unit. Additionally the underwater unit was equipped with a dissolved oxygen sensor and mounted in the bottom part of a Sea-Bird “SBE 12 Carousel Water Sampler” capable of holding 12 water-sampling bottles. The CTD oxygen data have not been validated onboard with water samples e.g. by iodometric oxygen titration. Therefore the oxygen sensor is considered unreliable.

# 6. Preliminary Results

## 6.1. Carbonates of the Gulf d'Arguin

H. Westphal, K. Gürs

Scientific targets

Carbonate production and deposition in high-nutrient tropical settings is rare in the modern world, and the northern part of Banc d'Arguin is probably the occurrence with the highest nutrient concentration. Elevated nutrients are caused not only by the upwelling system west of the Gulf d'Arguin, but also by high eolian influx of physically weathered dust. The carbonate deposits of Baie du Lévrier, the northernmost tip of the Banc d'Arguin have been described by Koopman et al. (1979), however a comprehensive study of the sediment-producing ecosystem was still lacking. In particular, this topic has high relevance as it can be expected that as a result of current global change, such high-nutrient, warm-water systems will become more common.

During Poseidon 346, within the shallow-water carbonate objectives we concentrated on (see Fig. 7):

- Surface sediment sampling in the Baie du Lévrier (grab sampling, box coring) for describing the modern sedimentary and ecological system;
- Surface sediment sampling along the 20-30 m isobath from Cap Blanc to Cap Timiris for studying the N-S trend from high-carbonate sediments (exceeding 90% carbonate contents) to sediments with as little as 40-50% carbonate;
- Vibro-coring of the coarse, pure carbonate sediments in order to gain information on environmental change through time.

Sampling was extremely successful, and a large data set was collected. One vibro-core was recovered in the Baie du Lévrier (Station 100-2) that is composed of coarse shell debris (Fig. 8). Because of the very shallow water depths (most parts have less than 10 m water depths), the vast shoals of the Banc d'Arguin could not be included in the sampling strategy. One sample was taken during a zodiac excursion on the southern shallow Banc d'Arguin, and one sample was taken by means of a bucket dredge.




Fig. 7: Sample locations on the Banc d'Arguin and its slopes.

## First results

### *Shallow-water carbonates of the Banc d'Arguin*

A clear trend in composition of the surface samples splits the Banc d'Arguin in three separate parts:

(1) The Baie du Lévrier and the northern part of the 20-30 m isobath section with clean, coarse carbonate sands;

(2) An area West and South of Cap Blanc, where clean quartzose sands are admixed that are interpreted as reworked dune sands;

(3) The southern part of the 20-30 m isobath section with large amounts of eolian silt admixed. These southern samples are overall fine-grained, poorly ventilated, black sediments;

(4) A fourth part is indicated by the dredge sample that consisted of skeletal sand with the only occurrence of red algal remains. However, to characterize this very shallow carbonate sedimentary realm, more systematic sampling would be required.




Fig. 8: Core GeoB 11601-02 (Station Pos 100-2) composed of coarse shell debris showing an overall coarsening-up trend.

Fig. 9 A: Coarse Fraction (> 2cm) from box core GeoB 11534-01 (Pos 346-34-1) from the northern part of Banc d'Arguin (water depth 29 m).




Fig. 9 B: Coarse Fraction (> 2cm) from grab sample GeoB 11596-01 (Pos 346-96-1) from the southern part of Banc d'Arguin (water depth 55 m). Dark elongated particles are worm tubes.

Fig. 9 C: Coarse Fraction (> 2cm) from grab sample GeoB 11508-01 (Pos 346-8-1) from the southern outer part of Banc d'Arguin (water depth 129 m). Abundant fish remains demonstrate high biological productivity.

Fig. 9 D: Coarse Fraction (> 2cm) from grab sample GeoB15597-01 (Pos 346-97-1) from the Baie du Lévrier (water depth 11 m). Sample is strongly monospecific with abundant *Capsella burnupi* (Sowerby, 1894).

The carbonate sedimentological assemblages reflect the high-nutrient situation of the study area. Zooxanthellate corals and green alga that are typical of oligotrophic tropical settings are absent from the samples collected. Generally, the carbonate fraction of the samples is dominated by mollusk remains (Fig. 9 A-D). The sediment is classified as heterozoan association sensu James (1997) or as foramol association sensu Lees and Buller (1972), clearly pointing out the atypical character of these tropical carbonates. The eutrophic character is manifested by high concentrations of fish remains (Fig. 9 C) including otoliths, and towards the south in the finer-grained deposits, by abundant worm tubes (Fig. 9 B).

From the carbonate grain composition of the samples, two areas are distinguished; (1) the northern part with the Baie du Lévrier and the northern part of the 20-30 m isobath section with abundant bivalves and distinctly less abundant gastropods (Fig. 9 A and D); (2) the southern part of the 20-30 m isobath section where worm tubes are dominant in many samples (Fig. 9 B).

In the Baie du Lévrier, some samples are highly monospecific with a high portion of the bivalve *Capsella burnupi* (Sowerby, 1894; Fig. 9 D and Fig. 10 A). This bivalve reflects the upwelling conditions in the study area. However, the sediment clearly has a warm-water, tropical character with the bivalve *Crassatina sp.* (Fig. 10 B) and gastropods like *Prunum annulatum* (Reeve, 1865) and *Marginella desjardini* (Marche-Marchad, 1957, Figs. 10 C and D) demonstrate that tropical water temperatures prevail.







A B

Fig. 10: Close-ups of environmentally significant mollusks (photographs by V. Diekamp, Marum, Bremen)

C D







*Sedimentation and faunal communities of the outer shelf and bathyal*

Some box cores and grab samples were taken on the outer shelf and bathyal slope West of the Banc d'Arguin and were complemented by grab samples West of Nouakchott (i.e. East of the Banda Mound Province). Aim was to examine the sedimentation outside the shallow inner shelf and the influx of upwelling on the faunal communities. A total of 12 box-cores and 18 grab samples were taken in water depth of below 50 m to 604 m.

West of the Banc d'Arguin all samples from the outer shelf consist of fine sand to silt interpreted as eolian dust from desert storms. Bathyal samples are composed of silts to clays and are of dark greenish grey colors. All samples are rich in shell debris. The outer shelf samples are dominated by bivalves of the family Telluride and the gastropod species *MacLean elate*. Deposit feeders of the Turritellidae (gastropods) are common. Similar associations are found for example in the Alboran sea of the western Mediterranean, where strong upwelling occurs, or in Middle Miocene deposits of the North Sea Basin (Janssen, 1984). Such associations indicate eutrophic conditions; for the North Sea Basin caused by freshwater influx (which for the Mauritania Shelf with its desert hinterland clearly is different). Despite high proliferation of organic matter no

signs of oxygen depletion are present in the samples. Exclusively tropical gastropods including *Clavatula sp*. or *Marginella spp*. confirm the tropical character of the sediments.

Stations Pos346-7 to Pos346-10 form a transect perpendicular to strike from shallow to deep water in the Southern part of the Gulf d'Arguin (Fig. 11):

The communities of Station Pos346-7 (Fig. 11 A) and Pos346-8 (Fig. 11 B) match with other outer shelf samples. Samples from Stations Pos346-7 to Pos346-9 contain relatively large amounts of allochthonous shallow water (intertidal) mollusk species mixed with the autochthonous fauna. Core Pos346-9-1 from a water depth of 939 m (GeoB11509-01; Fig. 11C) in addition contains rock fragments and deep-water corals of the species *Lophelia pertusa*; cf. Chapter 6.3). This erratic fauna is fundamentally different to the fauna described from the shores of the Banc d'Arguin (own unpublished data from samples from Tidra Island; Wolf et al., 1993). Thus it appears not to be transported to these sites from the East but could derive from rocky shoals known from the southern part of the Banc d'Arguin) and transported downhill by a northward directed countercurrent.

The community at Station Pos346-10 represents a typical bathyal community. Specialized carnivorous bivalves of the Cuspidariidae, deposit feeding bivalves of the Nuculanidae and the genus *Abra* as well as the deposit feeding gastropods including *Callumbonella suturalis* (Philippi, 1836) demonstrate the deep water character while no signs of anoxic or dysoxic conditions are present. In contrast, anoxic to dysoxic conditions are apparent in samples from Pos346-21-2 (GeoB11521-02; water depth 563 m) where Thyasiridae and *Lucinoma borealis* (Linnaeus, 1767) occur, both hosting chemosymbionts with the ability to oxidize sulfides (Fig. 11E).

The outer shelf off Nouakchott shows a contrasting picture. An isobathic section East of the Banda reefs in a water depth of about 150 m (Stations Pos346-71 to Pos346-75) reveals a strongly muddier facies where worms and Tellinidae are the dominant faunal elements. The fauna is diverse and particularly rich in deposit feeders. Solitary corals occur, and *Amyclina elata* (Gould, 1845) is virtually absent. In a transect perpendicular to strike towards the Banda reefs (Stations Pos346-75 to Pos346-77) the fauna is progressively impoverished downslope until the coarse fraction consists only of worm tubes, planktonic gastropods, and sparse shell fragments. This could imply stagnant water in the current shadow of the deep-water reefs.

In summary, the sediments from the outer shelf off the Banc d'Arguin reflect conditions influenced by strong upwelling. The environment is eutrophic, yet well oxygenized, and water temperatures are tropical. Conditions are well oxygenized in bathyal depth, whereas the outer shelf off Nouakchott, East of the Banda Mound Province, in contrast shows more stagnant eutrophic conditions. The upper bathyal of the slope in this area is nearly anoxic in the shadow of the deep-water reefs.

A




B

C

D

E

Fig. 11: Coarse fractions (> 2 cm) of samples from banc margin-slope transect at Southern Gulf d'Arguin. 0.5 cm squares on labels for scale. (C) Station Pos346-9: debris of deep-water coral *Lophelia pertusa*. (E) Station Pos346-21: bathyal community with chemosymbiontic affinity indicating dysoxic to anoxic conditions.

Further analytical work

Ongoing and planned analytical work includes

- Taxonomy of mollusks and other calcareous fauna
- Age dating by the radiocarbon method
- Grain size analysis
- Carbonate contents
- Stable isotope measurements on carbonate shells (incl. planktic foraminifers)

This work will lead to a comprehensive description of the ecological and depositional system of this high-nutrient tropical carbonate setting. This study will be the first of its kind systematically describing atypical tropical carbonates in terms of sedimentology, taxonomy, and geochemistry.

Study of the cores will add the time dimension, potentially revealing information on the development of upwelling in the past few thousand years. The study will also lead to a better understanding of the complex oceanographic situation with various water masses and strongly seasonal currents on this broad shallow-water area of the shelf of Mauritania. Integration with the siliciclastic data will add information on the climatic development in the hinterland. Thus, this study will add to the understanding of the coupled marine-terrestrial system of North-West Africa.

## 6.2. The Mud Belt of Banc d'Arguin

Till Hanebuth

Scientific targets

Two scientific targets were the focus of the mud belt objectives during Poseidon 346; the Late Holocene climate archive recorded in the mud belt (Fig. 12), and the question of the presence of a paleo-river draining into the Golf d'Arguin during the Holocene Climate Optimum.

*(a) Mud belt sediment of Banc d'Arguin as continuous archives for the reconstruction of the Late Holocene climate history*

The Banc d'Arguin is located off the Sahara desert. During the mid-Holocene Climate Optimum, the Sahara "Green mode" was characterized by a monsoonal precipitation rate several times higher then today and a comparably dense vegetation cover. Since then climatic deterioration has occurred in distinct steps with temporary recourses to more humid conditions until some two thousand years ago when modern hyper-arid conditions have finally established. This general development is known from different marine and terrestrial archives, however the evidence comes non-continuous or patchy records.

Mud belts that are usually located in certain physiographical positions in a sedimentary shelf system in many cases have the potential to continuously record the environmental history with very high temporal resolution. This potential is, however, rarely used up to now. The mud belt off Cap Timiris (in the following termed Timiris mud wedge) was first cored in 2005 (RV Meteor cruise M-65/2: Krastel et al., 2006) and current analyses (T. Hanebuth) give evidence that the Late Holocene is recorded in remarkable detail. Additionally, a second fine-grained wedge-like depocenter was discovered in an area around 20° N termed the Arguin mud wedge. (Fig. 12)

In order to expand the existing record back in time, during Poseidon 346 the seaward margin of both mud wedges along the mud belt was sampled. A series of surface samples taken around the mud belt was taken in order to study why the mud belt has such a stable position through time.




Fig. 12: Position of the mud belt consisting of the Arguin mud wedge and the Timiris mud wedge.

*(b) Question of presence of a paleo-river during the Holocene Climate Optimum draining into the Golf d'Arguin*

The existence of an extensive and deeply incised canyon system just off the Golf d'Arguin, together with the fluvial valley systems in the hinterland as visible on satellite images, and a number of endemic species from coastal waters, seem to indicate that a riverine supplier has been active in earlier, humid times. The Mid-Holocene green period of the Sahara would be a possible time interval of fluvial activity. However, this has not been proven to date because terrestrial and shelfal deposits disappear rapidly by microbial decay, widespread erosion, or are possibly overlain by thick sand seas and sediment sheets. Indirect and debatable evidence is given by marine proxies and is only related to grain size.

Subbottom structures on the outer shelf off the Golf d'Arguin, discovered during an earlier cruise (RV Meteor cruise M-65/2: Krastel et al., 2006) show erosional features in 80 m water depth that could be interpreted as of fluvial origin. Alternatively, they could represent a paleo-shoreline. During Poseidon 346, the material infilling these erosional structures was a prime target in order to answer the question of the origin of these sediment packages. In contrast to two earlier attempts employing a gravity corer, coring with the vibro-corer during Poseidon

346 resulted in complete recovery even in sediment succession with intercalated coarse-grained layers.

Results from the middle part of the Banc d'Arguin (TH)

*(a) Mud-belt depocenters*

Four sediment cores were taken from the Timiris mud wedge and 2 from the Arguin mud wedge. The sediments are composed of grayish homogenous silts that promise to represent a continuous succession that covers the past several thousand years. Core locations are placed at different positions at the margin of the depocenters, and dating will elucidate how these sediment bodies have evolved through time.

A zodiac trip towards one of the shoals at the margin of the Banc d'Arguin was undertaken to get an idea about the reasons why the Timiris mud wedge has developed just beyond a morphological step on the seafloor. A seafloor surface sample was composed of coarse-grained carbonate sand to gravel and supports the suggestion that a hydrological jump is responsible for massive grain-size separation in this area.

*(b) Paleo-shore vs. paleo-valley*

The five stations positioned on erosional features were chosen following shallow sediment acoustics from former research cruises. These cores are composed of a surficial sand facies overlaying fine to middle sands. These sands are interrupted by a massive shell bed showing grain-size sorting, good rounding and a faint fining-upward trend. Tentatively this shell horizon is interpreted as remnant of a paleo-coastline. Regarding the water depth of 80 m, they could be some 14 thousand years old – as a first preliminary estimate.




Fig. 13: Position of vibro-cores in the Timiris mud belt.




Fig. 14: Position of vibro-cores in the Timiris mud belt.




Fig. 15: Position of vibro-cores in the Arguin mud belt and the erosional features.




Fig. 16: Position of vibro-cores in the erosional features.

Next analytical steps

*(a) Analytics in the lab*

- Age dating by the radiocarbon method
- Grain-size analysis (sand, silt)
- Radiography
- XRF element scanning
- Stable isotope measurement on ostracods
- Component analysis
- Facies analysis

*(b) Scientific interpretation*

- Sedimentary facies interpretation with regard to the hydrological regime and coastline evolution (sea-level changes), respectively.
- Combination of resulting data with already existing core record from the *Timiris mud belt.*
- Comparison of mud belt data with terrestrial environmental records and archeological studies.
- Linkage of terrestrial history with shelf processes and depositional pattern on shelf and slope
- Facies model for the outer Golf d'Arguin in cooperation with the other cruise members by compilation of biological, calcareous and clastic surface data

## 6.3. Coral mounds

Markus Eisele and Karl Gürs

During Poseidon 346, two provinces of deep water carbonate mounds on the Mauritanian continental margin were surveyed: the *Banda Mound Province* (17°40'N, 16°40'W), named after the nearby Banda Gas Field and a newly described mound province that we named *Timiris Mound Province* (18°59'N, 16°52'W) after the close by Cape Timiris. The mounds were identified by low-resolution echo sounder profiles (Fig. 17) and sampled with the CTD unit, the Van Veen grab sampler, the box corer and the gravity corer. Grab samples, box cores and gravity cores were taken from the very tops of the mounds. Additionally one station (GeoB 11557) was located in the surrounding hemipelagic sediments near the mounds (off-mound) for reference.

The Banda Mound Province

The Banda mounds are located in water depths between 450 and 550 m. On the echo sounder, the Banda Mounds appear as conical build-ups with moats around their feet and thus show shapes typical for carbonate mounds covered by cold-water coral (Fig. 17). They reach heights of up to ~100 m above the surrounding sea floor.

Ten box cores and ten gravity cores were taken in the Banda Mound Province. The sediments are dominated by olive grey, clayey to silty, fine sands with varying contents of coral rubble and other bioclasts (Fig. 18 A and B). The fauna on the mounds shows a high biodiversity abundant corals (dominantly *Lophelia*, but *Madrepora* and *Dendrophyllia* are also present). Almost all corals retrieved were dead, with the exception of core GeoB 11564-1 where six living coral polyps (*Lophelia*) were found (Fig. 18 C.).

A conspicuous feature of the box cores is the high abundance of the bivalve *Acesta*. Other organisms typical for the Banda Mounds are ophiuroids, gorgonians, bryozoans, hydrozoans, sponges, crabs, squat lobsters, tube worms, and gastropods.




Fig. 17: Echo sounder profile of one of the Banda mounds

A 

B

C

Fig. 18: Examples of box cores from the Banda Mound province. (a) Upper surface, mud facies. (b) Upper surface, coral rubble facies with red gorgonian and a squat lobster. (c) Live polyps of *Lophelia*.

The Timiris Mound Province

The Timiris Mounds are located in depths similar to the Banda mounds further to the north. The echo sounder images of the Timiris Mounds again show typical patterns of cold-water coral covered carbonate mounds (Fig. 19). The mounds in this province also reach heights of some 100 m above the surrounding seafloor, comparable to those in the Banda Mound Province.




Fig. 19: Echo-sounder profile of a mound in the Timiris Mound Province.

Two box corers and two gravity cores were retrieved in the Timiris Mound Province. Sediments here are mainly composed of olive silty sand mixed with varying amounts of coral rubble (mainly *Lophelia* and abundant *Madrepora*) and other bioclasts (*Acesta*, agglutinated worm tubes, gastropods shells, and bryozoans) (Fig. 20). No live corals were found in the Timiris Mound Province. The fauna is composed of sponges, hydrozoans, bryozoans, snails, sea anemones and tube worms. One blade of seaweed was found in box core GeoB 11588-1.

A

B

C

Fig 20: Box cores from the Timiris Mound Province. (a) Surface of box core, sand-/ coral-rubble facies. (b) Surface of box core, sandy facies. (c) Seaweed in core GeoB 11588-1.

The faunal composition on the Mauritanian mounds differs pronouncedly from that of the coral mounds of the northern North Atlantic. For example the bivalve dominating the northern mounds, *Asperarca nodulosa* (Müller, 1776), and one of the most abundant gastropods *Amphissa acuticostata* (Philippi, 1844) are absent in the Mauritanian provinces.

A spectacular finding was that of some lazarus species including the gastropods *Cancellaria contorta* (Basterot, 1825) and *Narona lyrata* (Brocchi, 1814). These species are common in the Tertiary record of the North Atlantic and the Mediterranean and have vanished at the beginning of the Pleistocene. They apparently survived only in this unique habitat in the Mauritanian cold-water coral provinces.

## 6.4. CTD measurements

Jonas von Reumont

Introduction

A total of 34 CTD casts were performed during RV POSEIDON cruise 346 (Fig. 21). The measurements were conducted to generate profiles across deepwater coral mound structures from Cap Timiris to the Banda Mound region on the shelf slope off North Mauritania. Furthermore gradient parallel profiles in the Golf d'Arguin were carried out comprising part of the shelf and the shelf slope. The objective was to measure the characteristics of the water mass produced in the Golf d'Arguin by coastal upwelling and the Canary Current that is influencing the sedimentation of different carbonate sediments.




Figure 21: Overview of CTD stations in the study area.

Another objective was to study the stable isotope composition of oxygen and dissolved inorganic carbon as well as the strontium isotope composition in the water close to the sea floor on which the benthic organisms partly rely when building their skeletons. Where deeper casts were possible water samples were also taken within the water column.

Performance and Station Overview

The overall performance of the CTD was satisfactory. However, even though the recording computer worked without problems, some downcast measurements show numerous error measurement spikes within a water depth interval ranging from 10 to 100 m. The upcast measurements showed no systematic deviation from the downcast measurements but apart from individual spikes they did not replicate the error measurements and thus were chosen for further data processing. The data were processed using the Sea-Bird software "SBE Data Processing" Version 5.37d and visualized using "Ocean Data View" Version 3.2.0 (Schlitzer, R., http://odv.awi.de) (ftp://ftp.halcyon.com/pub/seabird/OUT/Seasoft-Win32/SBEDataProcessing).

The bottle release unit worked without problems, all bottles were closed reliably. During Station 63, sample bottle #5 was damaged and detached from the carousel. This accident might have been caused by the extraordinary high dolphin activity in the direct vicinity of the vessel. The bottle was replaced. Water samples were taken at all stations 10 m above seafloor (bottom

alarm) and at selected stations in the water column where significant changes in the measured parameters occurred.

In the Golf d'Arguin 22 CTD stations were measured. The profiles are positioned on the shelf and the shelf slope parallel to the gradient. A total of 12 CTD profiles have been measured in the region of coral mound areas. The transects were chosen perpendicular to the elongated mound structures. The water above the mounds was sampled as close to the sediment-water interface as possible.

Preliminary results

*Golf d'Arguin*

The profiles are oriented approximately perpendicular to the slope gradient. The general distribution of water masses is similar in all profiles within the Golf d'Arguin. Temperature and salinity show maximum values at the surface decreasing with depth. In conjunction with coastal upwelling in this region the northernmost profile shows clearly upward sloping isotherms towards the coast down to a water depth of 300 m (Fig. 22). Some profiles show a reversion of salinity and oxygen values within the first 200 m of the deepest casts (e.g. Fig. 22). Generally decreasing temperature and salinity but increasing dissolved oxygen below approx. 400 m of water depth (Fig. 23) may indicate the influence of a deeper current mentioned by Colman et al. (2005). Between 100 to 400 m oxygen values are lowest forming an oxygen minimum layer.




**Fig. 22**: Profiles of temperature (top), salinity (centre) and dissolved oxygen (bottom) comprising stations 30 to 34.




Fig. 23: Profiles of temperature (top), salinity (centre) and dissolved oxygen (bottom) comprising stations 50 to 54.




**Fig. 24: Surface plots of temperature (left), salinity (right) of the entire working area.**

Fig. 24 shows the distribution of sea surface temperature (SST) in the entire working area. In the north where upwelling water is expected to invade the Golf d'Arguin, SSTs are lowest, ranging from 17 to 19 °C. From 21°30.00 to 19°45.00 the measurements show an increase in temperature reaching 21 °C in the western segments of the profiles. Further south a continuous cooling of surface water down to 18 °C occurs. Salinity is highest in the north ranging from 36.3 to 36.0 psu with a general positive gradient from east to west. In contrast to temperature there is no increase in sea surface salinity between 21°30.00 to 19°45.00 but a continuous overall decrease from north to south down to 35.8 psu.

The T-S diagram of the Golf d'Arguin region (Fig. 25) shows the meridional changes in surface and near surface waters as a considerable spreading of data values. Below 800 m of water depth the diagram seems to indicate the admixture of a colder and more saline water mass.




Fig. 25: T-S diagram of CTD measurements in the Golf d'Arguin region.

*Carbonate mound region*

12 CTD casts were conducted at the location or in the vicinity of carbonate mound structures (Banda Mound and Timiris Mound provinces). The measurements of temperature and salinity show a horizontally well-stratified water column with highest values at the surface and a decrease with depth. Compared to the Golf d'Arguin, a more pronounced thermocline is developed in the mound regions (Figs. 26 and 27) representing less mixing of the water column. Furthermore the oxygen gradient in the upper 100 m is slightly steeper and the minimum zone extends to a greater depth. Below 600 m of water depth a pronounced increase of oxygen values is observed. Apparently the oxygen minimum layer is also represented by the bending of the curve in the T-S diagram of the carbonate mound region (Fig. 28) that roughly correlates with the depth of the upper part of this layer.

Except for slightly higher temperature and salinity values above the mound top compared to lateral values there are now local anomalies associated with the carbonate mound structures.




Fig. 26: Profiles of temperature (top), salinity (centre) and dissolved oxygen (bottom) comprising stations 82 to 86b.




Fig. 27: Profiles of temperature (top), salinity (centre) and dissolved oxygen (bottom) comprising stations 59 to 61.




Fig. 28: T-S diagram of CTD measurements in the carbonate mound area south of Cape Timiris.

# 7. Weather and Meteorological Conditions

Generally, the conditions during the cruise were calm with low to no waves, which was particularly important for operating the vibro-corer. However, storms (Jan. 2nd; 11th) interrupted the work process, and we had to terminate the work on 11. of Jan. and thus 24h earlier than planned, and returned to Las Palmas. These storms were attributed to the Harmattan, a Northeast trade wind. The effect of these winter storms on the Saharan hinterland was apparent even under calm weather conditions by extreme dust loading of the air, for several days restricting visibility to less than 1 mile.
Generally, the sea was rougher in the mornings and calmed down around noon. The afternoons on many days were entirely lacking any waves.
On Jan. 10th and 11th, we experienced humid and rainy weather.

Fig. 29: Landsat image from January 1st, 2007 illustrating the dust storm R/V Poseidon experienced. (http://rapidfire.sci.gsfc.nasa.gov/gallery/?2007001-0101/WestAfrica.A2007001.1455.1km.jpg)

## 8. Acknowledgements

The Scientific Shipboard Party aboard RV Poseidon during POS 346 gratefully acknowledges the friendly cooperation and very efficient technical support of Captain Michael Schneider, his officers and crew who substantially contributed to the overall scientific success of this cruise. We also appreciate the great help of the Leitstelle in Kiel, namely Tom Müller. RCOM supported this cruise technically and logistically (Götz Ruhland and others). The work was funded by the German Science Foundation through project We 2493-MACUYS, HERMES, and ship time to HW and AF.

## 9. References

Colman, J.G., Gordon, D.M., Lane, A.P., Forde, M.J, Fitzpatrick, J.J. (2005): Carbonate mounds off Mauritania, Northwest Africa: status of deep-water corals and implications for management of fishing and oil exploration activities. In: Freiwald, A. and Roberts, J.M. (eds.): Cold-Water Corals and Ecosystems, Erlangen Earth Conference Series. Springer, Berlin, Heidelberg.

James, N.P. (1997) The cool-water carbonate depositional realm. In: Cool-Water Carbonates (Eds N.P. James and J. Clarke), SEPM Spec. Publ., 56, 1–20.

Janssen, A.W. (1984) Mollusken uit het Miozeen van Winterswijk-Miste, Bibliotheek van de Koninklijk Nederlandse Natuurhistorische Vereniging, 36, 451 pp

Koopman, B. J., Lees, A., Piessens, P., Sarnthein, M., 1979, Skeletal carbonate sands and wind-derived silty marls off the Saharan coast: Baie du Lévrier, Aguin Platform, Mauritania. "Meteor" Forschungs-Ergebnisse C, 30, 15-57.

Krastel, S., Hanebuth, T.J.J., et al., 2006. Report and preliminary results of RV Meteor cruise M65/2 Dakar. 767, 185, Universität Bremen, Bremen

Lees, A. and Buller, A.T. (1972) Modern temperate water and warm water shelf carbonate sediments contrasted. Mar. Geol., 13, 67–73.

Wolf, W. J., A. G. Duiven, P . Duiven, P . Esselink, Abou Gueye, A. Meijboom, G. Moerland & J . Zegers (1993) Biomass of macrobenthic tidal flat fauna of the Banc d'Arguin, Mauritania. Hydrobiologia 258 : 151-163, Kluwer, Belgium.

Appendix B: Stations list

| Station number | GeoB-number | Date | Time begin of station | Latitude | Longitude | Water depth | Tool |
|---|---|---|---|---|---|---|---|
| POS 346-01 | 115-01 | 30.12.2006 | 21:43 | 19°45.0000 | 017°29.9900 | 1550,0 | CTD |
| POS 346-02-1 | 115-02-1 | 31.12.2006 | 8:15 | 19°40.2468 | 016°57.9678 | 58,0 | GBC |
| POS 346-02-2 | 115-02-2 | 31.12.2006 | 8:15 | 19°40.2540 | 016°57.9372 | 58,0 | GC |
| POS 346-03-1 | 115-03-1 | 31.12.2006 | 9:25 | 19°41.0448 | 016°59.1438 | 67,0 | GBC |
| POS 346-03-2 | 115-03-2 | 31.12.2006 | 9:25 | 19°41.0442 | 016°59.1480 | 67,0 | GC |
| POS 346-04-1 | 115-04-1 | 31.12.2006 | 10:16 | 19°39.7050 | 016°59.1348 | 75,0 | GBC |
| POS 346-04-2 | 115-04-2 | 31.12.2006 | 10:16 | 19°39.6810 | 016°59.1252 | 74,0 | GC |
| POS 346-05-1 | 115-05-1 | 31.12.2006 | 11:21 | 19°39.6648 | 016°58.2360 | 64,0 | GBC |
| POS 346-05-2 | 115-05-2 | 31.12.2006 | 11:21 | 19°39.6612 | 016°58.2210 | 64,0 | GC |
| POS 346-46-1 | 115-46-1 | 31.12.2006 | 13:07 | 19°39.7662 | 017°00.6372 | 104,0 | VVG |
| POS 346-46-2 | 115-46-2 | 31.12.2006 | 13:07 | 19°39.7422 | 017°00.6030 | 103,0 | VVG |
| POS 346-06 | 115-06 | 31.-01.01.2007 | | | | | ESR |
| POS 346-07-1 | 115-07-1 | 01.01.2007 | 9:55 | 19°44.4888 | 017°03.0360 | 73,0 | VVG |
| POS 346-07-2 | 115-07-2 | 01.01.2007 | 9:55 | 19°44.4918 | 017°03.0228 | 73,0 | GBC |
| POS 346-07-3 | 115-07-3 | 01.01.2007 | 9:55 | 19°44.4858 | 017°03.0282 | 73,0 | GC |
| POS 346-08-1 | 115-08-1 | 01.01.2007 | 11:10 | 19°43.5000 | 017°05.9952 | 129,0 | VVG |
| POS 346-08-2 | 115-08-2 | 01.01.2007 | 11:10 | 19°43.5102 | 017°05.9700 | 128,0 | GBC |
| POS 346-09-1 | 115-09-1 | 01.01.2007 | 12:55 | 19°42.4938 | 017°42.4938 | 370,0 | GBC |
| POS 346-10-1 | 115-10-1 | 01.01.2007 | 14:06 | 19°41.5572 | 017°12.0222 | 600,0 | GBC |
| POS 346-10-2 | 115-10-2 | 01.01.2007 | 14:06 | 19°41.5752 | 017°11.9982 | 598,0 | GC |
| POS 346-11 | 115-11 | 01.01.2007 | 16:00 | 19°42.5718 | 017°09.3462 | 454,0 | VVG |
| POS 346-11A | 115-11A | 01.01.2007 | 17:49 | 19°46.9908 | 017°03.9888 | 22,7 | VVG |
| POS 346-11B | 115-11B | 01.01.2007 | 19:10 | 19°50.5398 | 017°11.9580 | 24,2 | VVG |
| POS 346-11C | 115-11C | 01.01.2007 | 19:58 | 19°55.0290 | 017°12.4440 | 27,1 | VVG |
| POS 346-11D | 115-11D | 01.01.2007 | 20:57 | 20°00.0222 | 017°14.9952 | 26,1 | VVG |
| POS 346-11E | 115-11E | 01.01.2007 | 21:56 | 20°19.0302 | 017°19.0302 | 28,3 | VVG |
| POS 346-11F | 115-11F | 01.01.2007 | 22:41 | 20°06.3720 | 017°21.0618 | 27,4 | VVG |
| POS 346-11G | 115-11G | 01.01.2007 | 23:19 | 20°09.0162 | 017°22.5420 | 27,1 | VVG |
| POS 346-13-1 | 115-13-1 | 02.01.2007 | 12:46 | 20°33.0080 | 017°12.0408 | 31,3 | VVG |
| POS 346-13-2 | 115-13-2 | 02.01.2007 | 12:46 | 20°33.9852 | 017°12.0300 | 31,2 | GBC |
| POS 346-14-1 | 115-14-1 | 02.01.2007 | 14:24 | 20°38.0130 | 017°07.0428 | 19,0 | GBC |
| POS 346-15-1 | 115-15-1 | 02.01.2007 | 15:41 | 20°40.9998 | 017°02.0262 | 16,8 | GBC |
| POS 346-15-2 | 115-15-2 | 02.01.2007 | 15:41 | 20°40.9728 | 017°02.0160 | 16,1 | VL |
| POS 346-16-1 | 115-16-1 | 02.01.2007 | 17:12 | 20°43.9980 | 016°57.2010 | 9,1 | GBC |
| POS 346-17-1 | 115-17-1 | 02.01.2007 | 18:10 | 20°41.9650 | 017°02.0310 | 17,3 | VL |
| POS 346-18 | 115-18 | 02.01.2007 | 22:12 | 20°24.05 | 017°23.94 | 39,1 | CTD |
| POS 346-19 | 115-19 | 02.01.2007 | 23:28 | 20°23.90 | 017°31.91 | 58,0 | CTD |
| POS 346-20 | 115-20 | 03.01.2007 | 0:46 | 20°24.43 | 017°40.98 | 143,0 | CTD |
| POS 346-21-1 | 115-21-1 | 03.01.2007 | 2:00 | 20°23.9070 | 017°47.9562 | 559,0 | CTD |
| POS 346-21-2 | 115-21-2 | 03.01.2007 | 7:59 | 20°24.0612 | 017°48.0108 | 560,0 | GBC |
| POS 346-21-3 | 115-21-3 | 03.01.2007 | 7:59 | 20°24.0102 | 017°47.9988 | 557,0 | GC |
| POS 346-22-1 | 115-22-1 | 03.01.2007 | 10:22 | 20°24.5262 | 017°41.0058 | 151,0 | GBC |
| POS 346-23-1 | 115-23-1 | 03.01.2007 | 11:57 | 20°23.9712 | 017°31.9950 | 59,0 | GBC |
| POS 346-24-1 | 115-24-1 | 03.01.2007 | 13:16 | 20°23.9820 | 017°23.9478 | 38,5 | VVG |
| POS 346-24-2 | 115-24-2 | 03.01.2007 | 13:16 | 20°23.9220 | 017°23.9178 | 38,4 | GBC |
| POS 346-24-3 | 115-24-3 | 03.01.2007 | 13:16 | 20°23.8938 | 017°23.9028 | 39,1 | GBC |
| POS 346-25-1 | 115-25-1 | 03.01.2007 | 15:28 | 20°29.7528 | 017°16.5438 | 34,1 | VVG |
| POS 346-25-2 | 115-25-2 | 03.01.2007 | 15:28 | 20°29.6598 | 017°16.5480 | 33,1 | GBC |
| POS 346-13-3 | 115-13-1 | 03.01.2007 | 16:48 | 20°33.9360 | 017°12.0432 | 31,4 | VL |
| POS 346-26-1 | 115-26-1 | 03.01.2007 | 19:16 | 20°42.4620 | 017°07.9838 | 25,2 | VVG |
| POS 346-27-1 | 115-27-1 | 03.01.2007 | 19:58 | 20°41.9940 | 017°04.9740 | 15,8 | VVG |
| POS 346-28-1 | 115-28-1 | 03.01.2007 | 20:45 | 20°38.0058 | 017°01.9778 | 18,0 | VVG |
| POS 346-29-1 | 115-29-1 | 03.01.2007 | 21:35 | 20°36.5202 | 017°04.9578 | 19,0 | VVG |
| POS 346-30-1 | 115-30-1 | 03.01.2007 | 22:27 | 20°33.5232 | 017°09.0120 | 28,9 | VVG |
| POS 346-31-1 | 115-31-1 | 04.01.2007 | 0:13 | 20°27.4500 | 017°14.4288 | 30,8 | VVG |
| POS 346-32-1 | 115-32-1 | 04.01.2007 | 1:00 | 20°23.4498 | 017°14.5668 | 31,0 | VVG |
| POS 346-33-1 | 115-33-1 | 04.01.2007 | 2:06 | 20°17.9448 | 017°19.0452 | 34,0 | VVG |
| POS 346-34-1 | 115-34-1 | 04.01.2007 | 2:49 | 20°19.4220 | 017°21.0780 | 29,0 | VVG |
| POS 346-35-1 | 115-35-1 | 04.01.2007 | 3:44 | 20°13.9302 | 017°22.5582 | 32,0 | VVG |
| POS 346-36-1 | 115-36-1 | 04.01.2007 | 8:04 | 20°03.5808 | 017°28.9020 | 54,0 | GBC |
| POS 346-37-1 | 115-37-1 | 04.01.2007 | 8:50 | 20°05.4852 | 017°32.5812 | 78,0 | VVG |
| POS 346-38-1 | 115-38-1 | 04.01.2007 | 9:17 | 20°05.0628 | 017°33.6612 | 86,0 | VVG |
| POS 346-39-1 | 115-39-1 | 04.01.2007 | 9:35 | 20°04.8828 | 017°34.0332 | 88,0 | GBC |
| POS 346-40-1 | 115-40-1 | 04.01.2007 | 10:09 | 20°04.3722 | 017°33.1650 | 85,0 | VVG |
| POS 346-41-1 | 115-41-1 | 04.01.2007 | 10:29 | 20°03.7878 | 017°33.2748 | 85,0 | VVG |
| POS 346-41A-1 | 115-41A-1 | 04.01.2007 | 11:10 | 20°01.3932 | 017°34.3530 | 171,0 | VVG |
| POS 346-41A-2 | 115-41A-2 | 05.01.2007 | 11:10 | 20°01.3728 | 017°34.3542 | 178,0 | VVG |
| POS 346-41B | 115-41B | 04.01.2007 | 12:01 | 20°01.5000 | 017°31.4718 | 75,0 | VVG |
| POS 346-41C | 115-41C | 04.01.2007 | 12:32 | 20°03.0300 | 017°33.8118 | 89,0 | VVG |

Appendix B: Stations list

| Station number | GeoB-number | Date | Time begin of station | Latitude | Longitude | Water depth | Tool |
|---|---|---|---|---|---|---|---|
| POS 346-41D | 115-41D | 04.01.2007 | 13:26 | 20°00.6888 | 017°36.1332 | 329,0 | VVG |
| POS 346-41E | 115-41E | 04.01.2007 | 14:14 | 20°00.8838 | 017°38.1990 | 502,0 | VVG |
| POS 346-36-2 | 115-36-2 | 04.01.2007 | 16:02 | 20°03.5850 | 017°28.9500 | 54,0 | GC |
| POS 346-38-2 | 115-38-2 | 04.01.2007 | 16:53 | 20°05.0910 | 017°33.6780 | 86,0 | VL |
| POS 346-39-2 | 115-39-2 | 04.01.2007 | 17:55 | 20°04.9212 | 017°33.9702 | 87,0 | VL |
| POS 346-40-2 | 115-40-2 | 04.01.2007 | 18:53 | 20°04.4058 | 017°33.1392 | 85,0 | VL |
| POS 346-42-1 | 115-42-1 | 04.01.2007 | 23:56 | 19°56.4600 | 017°42.4200 | 296,0 | CTD |
| POS 346-43-1 | 115-43-1 | 05.01.2007 | 1:19 | 19°59.8800 | 017°36.9900 | 624,0 | CTD |
| POS 346-44-1 | 115-44-1 | 05.01.2007 | 3:13 | 20°03.9552 | 017°31.0362 | 69,0 | CTD |
| POS 346-45-1 | 115-45-1 | 05.01.2007 | 4:33 | 20°07.9692 | 017°25.0038 | 31,0 | CTD |
| POS 346-47-1 | 115-47-1 | 05.01.2007 | 8:02 | 20°25.5600 | 017°12.9802 | 25,6 | BG |
| POS 346-48-1 | 115-48-1 | 05.01.2007 | 9:23 | 20°20.5272 | 017°16.9858 | 28,0 | BG |
| POS 346-49-1 | 115-49-1 | 05.01.2007 | 10:35 | 20°17.0238 | 017°21.6708 | 28,9 | BG |
| POS 346-36-3 | 115-36-3 | 05.01.2007 | 15:12 | 20°03.6700 | 017°29.0300 | 54,0 | VL |
| POS 346-99-1 | 115-99-1 | 05.01.2007 | 16:22 | 20°04.7500 | 017°33.1700 | 85,0 | VL |
| POS 346-41-2 | 115-41-2 | 05.01.2007 | 17:11 | 20°03.9390 | 017°33.2190 | 85,0 | VL |
| POS 346-50 | 115-50 | 05.01.2007 | 22:10 | 19°41.5500 | 017°11.9100 | 597,0 | CTD |
| POS 346-51 | 115-51 | 05.01.2007 | 23:07 | 19°42.5500 | 017°08.9400 | 424,0 | CTD |
| POS 346-52 | 115-52 | 05.01.2007 | 23:55 | 19°43.5500 | 017°05.9000 | 128,0 | CTD |
| POS 346-53 | 115-53 | 06.01.2007 | 0:36 | 19°43.4900 | 017°03.0100 | 90,0 | CTD |
| POS 346-54 | 115-54 | 06.01.2007 | 1:10 | 19°44.4900 | 017°03.0000 | 73,0 | CTD |
| POS 346-55 | 115-55 | 06.01.2007 | | | | | ESR |
| POS 346-56 | 115-56 | 06.01.2007 | | | | | ESR |
| POS 346-58 | 115-58 | 06.01.2007 | 20:15 | 17°41.4000 | 016°39.9270 | 482,0 | CTD |
| POS 346-59 | 115-59 | 06.01.2007 | 20:54 | 17°41.0009 | 016°40.1988 | 544,0 | CTD |
| POS 346-60 | 115-60 | 06.01.2007 | 21:25 | 17°41.0052 | 016°40.1070 | 480,0 | CTD |
| POS 346-61 | 115-61 | 06.01.2007 | 21:57 | 17°41.1000 | 016°39.9222 | 548,0 | CTD |
| POS 346-62 | 115-62 | 06.01.2007 | 23:31 | 17°39.9700 | 016°33.0500 | 195,0 | CTD |
| POS 346-63 | 115-63 | 07.01.2007 | 1:40 | 17°36.1400 | 016°47.9700 | 1030,0 | CTD |
| POS 346-64-1 | 115-64-1 | 07.01.2007 | 7:57 | 17°38.7510 | 016°39.9888 | 467,0 | GBC |
| POS 346-64-2 | 115-64-2 | 07.01.2007 | 7:57 | 17°38.7528 | 016°39.9552 | 442,0 | GC |
| POS 346-65-1 | 115-65-1 | 07.01.2007 | 9:29 | 17°38.9928 | 016°39.7512 | 456,0 | GBC |
| POS 346-65-2 | 115-65-2 | 07.01.2007 | 9:29 | 17°39.0000 | 016°39.7380 | 440,0 | GC |
| POS 346-66-1 | 115-66-1 | 07.01.2007 | 10:41 | 17°39.0078 | 016°40.0842 | 441,0 | GBC |
| POS 346-66-2 | 115-66-2 | 07.01.2007 | 10:41 | 17°39.0000 | 016°40.0722 | 441,0 | GC |
| POS 346-67-1 | 115-67-1 | 07.01.2007 | 13:00 | 17°39.5100 | 016°40.0668 | 428,0 | GBC |
| POS 346-67-2 | 115-67-2 | 07.01.2007 | 13:00 | 17°39.4968 | 016°40.0338 | 429,0 | GC |
| POS 346-57-1 | 115-57-1 | 07.01.2007 | 14:22 | 17°41.2458 | 016°39.7392 | 478,0 | GBC |
| POS 346-57-2 | 115-57-2 | 07.01.2007 | 14:22 | 17°41.2470 | 016°39.7340 | 477,0 | GC |
| POS 346-70-1 | 115-70-1 | 07.01.2007 | 23:56 | 17°56.9760 | 016°28.9080 | 158,0 | BG |
| POS 346-71-1 | 115-71-1 | 08.01.2007 | 0:52 | 17°52.9728 | 016°28.9542 | 157,0 | BG |
| POS 346-72-1 | 115-72-1 | 08.01.2007 | 1:47 | 17°48.0120 | 016°29.0100 | 153,0 | BG |
| POS 346-73-1 | 115-73-1 | 08.01.2007 | 2:39 | 17°43.9842 | 016°29.0352 | 148,0 | BG |
| POS 346-74-1 | 115-74-1 | 08.01.2007 | 3:23 | 17°39.9888 | 016°28.9890 | 138,0 | VVG |
| POS 346-75-1 | 115-75-1 | 08.01.2007 | 4:00 | 17°40.0182 | 016°31.0050 | 167,0 | VVG |
| POS 346-76-1 | 115-76-1 | 08.01.2007 | 4:39 | 17°39.9912 | 016°33.0450 | 194,0 | VVG |
| POS 346-77-1 | 115-77-1 | 08.01.2007 | 5:12 | 17°39.9972 | 016°35.0460 | 246,0 | VVG |
| POS 346-68-1 | 115-68-1 | 08.01.2007 | 8:14 | 17°39.8562 | 016°40.3542 | 514,0 | GBC |
| POS 346-68-2 | 115-68-2 | 08.01.2007 | 8:14 | 17°39.8550 | 016°40.3608 | 493,0 | GC |
| POS 346-69-1 | 115-69-1 | 08.01.2007 | 9:30 | 17°40.0080 | 016°40.3230 | 440,0 | GBC |
| POS 346-69-2 | 115-69-2 | 08.01.2007 | 9:30 | 17°40.0062 | 016°40.3272 | 444,0 | GC |
| POS 346-78-1 | 115-78-1 | 08.01.2007 | 10:30 | 17°40.5132 | 016°40.2012 | 458,0 | GBC |
| POS 346-78-2 | 115-78-2 | 08.01.2007 | 10:30 | 17°40.5162 | 016°40.2060 | 453,0 | GC |
| POS 346-79-1 | 115-79-1 | 08.01.2007 | 12:51 | 17°40.7658 | 016°40.0842 | 452,0 | GBC |
| POS 346-79-2 | 115-79-2 | 08.01.2007 | 12:51 | 17°40.7628 | 016°40.1040 | 450,0 | GBC |
| POS 346-79-3 | 115-79-3 | 08.01.2007 | 12:51 | 17°40.7688 | 016°40.0980 | 451,0 | GC |
| POS 346-80-1 | 115-80-1 | 08.01.2007 | 14:28 | 17°41.0388 | 016°40.0782 | 481,0 | GBC |
| POS 346-80-2 | 115-80-2 | 08.01.2007 | 14:28 | 17°40.9932 | 016°40.0662 | 483,0 | GC |
| POS 346-81-1 | 115-81-1 | | | | | | ESR |
| POS 346-82-1 | 115-82-1 | 09.01.2007 | 2:02 | 18°58.9800 | 016°54.4000 | 734,0 | CTD |
| POS 346-83-1 | 115-83-1 | 09.01.2007 | 3:08 | 18°59.0300 | 016°53.1900 | 643,0 | CTD |
| POS 346-84-1 | 115-84-1 | 09.01.2007 | 4:03 | 18°58.9800 | 016°52.0700 | 566,0 | CTD |
| POS 346-85-1 | 115-85-1 | 09.01.2007 | 4:39 | 18°58.9900 | 016°51.9600 | 483,0 | CTD |
| POS 346-86A-1 | 115-86A-1 | 09.01.2007 | 5:10 | 18°58.9800 | 016°51.7800 | 465,0 | CTD |
| POS 346-86B-1 | 115-86B-1 | 09.01.2007 | 5:44 | 18°58.9700 | 016°51.5500 | 536,0 | CTD |
| POS 346-88-1 | 115-88-1 | 09.01.2007 | 8:01 | 18°58.9920 | 016°51.8178 | 474,0 | GBC |
| POS 346-87-1 | 115-87-1 | 09.01.2007 | 8:42 | 18°58.9950 | 016°51.9348 | 482,0 | GBC |
| POS 346-87-2 | 115-87-2 | 09.01.2007 | 8:42 | 18°58.9938 | 016°51.9192 | 482,0 | GC |
| POS 346-89-1 | 115-89-1 | 09.01.2007 | 10:52 | 19°06.3948 | 016°50.5050 | 860,0 | CTD |

Appendix B: Stations list

| Station number | GeoB-number | Date | Time begin of station | Latitude | Longitude | Water depth | Tool |
|---|---|---|---|---|---|---|---|
| POS 346-90-1 | 115-90-1 | 09.01.2007 | 15:15 | 19°13.0020 | 016°37.6902 | 65,0 | CTD |
| POS 346-91A | 115-91A | 09.01.2007 | 17:15 | 19°44.1900 | 016°04.7920 | 3.5 to 5 | BD |
| POS 346-91B | 115-91B | 09.01.2007 | 17:15 | 19°44.1900 | 016°04.7920 | 3.5 to 5 | BD |
| POS 346-91C | 115-91C | 09.01.2007 | 17:15 | 19°44.1900 | 016°04.7920 | 3.5 to 5 | BD |
| POS 346-92-1 | 115-92-1 | 09.01.2007 | 20:40 | 19°37.0000 | 016°53.7900 | 33,8 | CTD |
| POS 346-93-1 | 115-93-1 | 09.01.2007 | 21:39 | 19°43.4868 | 016°57.2748 | 23,3 | VVG |
| POS 346-94-1 | 115-94-1 | 09.01.2007 | 22:15 | 19°44.4888 | 016°59.9748 | 33,0 | VVG |
| POS 346-95-1 | 115-95-1 | 09.01.2007 | 22:54 | 19°45.8100 | 017°02.0040 | 37,2 | VVG |
| POS 346-96-1 | 115-96-1 | 09.01.2007 | 23:30 | 19°47.2938 | 017°05.7750 | 55,0 | VVG |
| POS 346-97-1 | 115-97-1 | 10.01.2007 | 8:39 | 20°42.7338 | 017°03.5328 | 11,1 | VVG |
| POS 346-97-2 | 115-97-2 | 10.01.2007 | 8:39 | 20°42.7418 | 017°03.5438 | 11,3 | VVG |
| POS 346-100-1 | 116-01-1 | 10.01.2007 | 9:30 | 20°42.4992 | 016°59.0112 | 12,2 | VVG |
| POS 346-100-2 | 116-01-2 | 10.01.2007 | 9:30 | 20°42.4920 | 016°58.9620 | 12,1 | VL |
| POS 346-101-1 | 116-02-1 | 10.01.2007 | 11:11 | 20°38.1638 | 017°03.9642 | 17,6 | VVG |
| POS 346-102-1 | 116-03-1 | 10.01.2007 | 12:05 | 20°42.2238 | 017°02.7670 | 20,2 | VVG |
| POS 346-103-1 | 116-04-1 | 10.01.2007 | 13:31 | 20°48.9042 | 017°00.9960 | 11,8 | VVG |
| POS 346-104-1 | 116-05-1 | 10.01.2007 | 15:13 | 20°45.1812 | 017°03.4638 | 11,0 | VVG |
| POS 346-104-2 | 116-05-2 | 10.01.2007 | 15:13 | 20°45.1812 | 017°03.4638 | 12,1 | VVG |
| POS 346-104-3 | 116-05-3 | 10.01.2007 | 15:13 | 20°45.2010 | 017°03.4632 | 11,0 | VL |
| POS 346-98-1 | 116-06-1 | 10.01.2007 | 16:47 | 20°43.1730 | 017°02.7930 | 20,1 | VVG |
| POS 346-105-1 | 116-07-1 | 10.01.2007 | 17:30 | 20°45.4858 | 016°59.1000 | 13,6 | VVG |
| POS 346-106-1 | 116-08-1 | 10.01.2007 | 23:59 | 20°41.9400 | 017°21.5000 | 52,0 | CTD |
| POS 346-107-1 | 116-09-1 | 11.01.2007 | 1:14 | 20°42.0600 | 017°29.9300 | 67,0 | CTD |
| POS 346-108-1 | 116-10-1 | 11.01.2007 | 2:38 | 20°41.9800 | 017°38.0800 | 97,0 | CTD |
| POS 346-109-1 | 116-11-1 | 11.01.2007 | 3:15 | 20°42.1000 | 017°42.4100 | 389,0 | CTD |
| POS 346-110-1 | 116-12-1 | 11.01.2007 | 4:26 | 20°42.0300 | °46.7500+F35 | 498,0 | CTD |
| POS 346-111-1 | 116-13-1 | 11.01.2007 | 8:10 | 20°41.9780 | 017°38.1918 | 98,0 | VVG |
| POS 346-111-2 | 116-13-2 | 11.01.2007 | 8:10 | 20°42.0030 | 017°38.1930 | 99,0 | GBC |
| POS 346-111-3 | 116-13-3 | 11.01.2007 | 8:10 | 20°42.8228 | 017°38.1848 | 100,0 | VL |
| POS 346-112-1 | 116-14-1 | 11.01.2007 | 10:50 | 20°42.9550 | 017°29.9958 | 68,0 | VVG |
| POS 346-112-2 | 116-14-2 | 11.01.2007 | 10:50 | 20°42.9550 | 017°29.9958 | 67,0 | GBC |

GBC=giant box corer; GC=Gravity corer; VL=vibrocorer; VVG=Van Veen grab; BD=bucked dredge (dinghy); ESR=echo sounder reconnaissance

Poseidon 346 MACUMA
1. Wochenbericht
28.12.2006-3.1.2007

Die MACUMA Forschungsfahrt auf FS Posiedon befaßt sich mit dem gemischt karbonatisch-siliziklastischen Ablagerungssystemen auf dem Schelf und Schelfhang von NW-Afrika vor Mauretanien (Banc d'Arguin und westlich Nouakchott). Das Arbeitsgebiet liegt imTropengürtel (17-21°N), jedoch im Einfluß von Kanarenstrom und einer Upwelling-Front, so daß sich kein typisch tropisches Karbonatablagerungssystem mit zooxanthellaten Riffen ausgebildet hat. Stattdessen kommen atypische tropische Karbonate ohne zooxanthellate Korallen und Grünalgen vor. Auf dem Schelfhang vor Mauretanien befinden sich als weiteres Element des Ökosystems wenig bekannte Tiefwasserriffe. Weiterhin ist das gemischt kabonatisch-siliziklastische System für die Klimarekonstruktion des Quartärs von Bedeutung, wie Vorarbeiten an einem Kern von der südlichen Banc d'Arguin zeigen.

Die Fragestellungen für diese Ausfahrt umfassen:

- Erfassung und Beschreibung dieses rezenten atypischen tropischen Karbonatablagerungssystems (Sedimentologie, Taxonomie, Geochemie, Ozeanographie)
- Untersuchung der Tiefwasserriffe am Kontinentalhang
- Untersuchung der quartären Klimaentwicklung anhand von (fluviatilen und äolischen) siliziklastischen sowie karbonatischen Sedimentkomponenten.

28.12.2006
Auslaufen um 9:00; Transit zur ersten Station (knapp 62 h Transit-Zeit).

30.12.2006
Station 1 um 22:45: CTD (vor der Schelfkante, mittlere Banc d'Arguin; 1550 m). Das Temperatur- und das Salinitätsprofil zeigen deutlich den nur 50 m mächtigen Kanarenstrom.

31.12.2006
Stationen 2-5, südliche Banc d'Arguin. Lage in einem Mudbelt. Es wurden GKG und SL gefahren, der Kerngewinn war sehr gut. Das Material ist dunkel, siltig und relativ reich an Mollusken. Eine zusätzliche Station wurde westlich des Mudbelt gefahren, um das Vibrolot auszuprobieren. Der vorweggeschickte Backengreifer war allerdings leer. Das Vibrolot hatte eine technische Panne, so daß die Station abgebrochen wurde. Um 15:00 war Arbeitsende wegen Feiertages. Nachts wurde ein Echolotprofil gefahren (Station 6).

1.1.2007
Stationen 7-10 bilden ein Profil senkrecht zur Schelfkante nördlich des Mudbelt auf der südlichen Banc d'Arguin. Stationen 7 und 8 wurden mit Backengreifer und mit GKG beprobt. Stationen 9 und 10 wurden mit GKG beprobt. Das Sediment ist anoxisch und riecht nach $H_2S$. Die Fauna spiegelt nährstoffreiche Bedingungen wieder. Mehrere Kerne brachten lebende Fische mit an Deck. Der GKG von Station 9 enthält Solitärkorallen und Bruchstücke von *Madrepora*. Eine neue Station 11 wurde festgelegt, die auf einer Erhebung im Echolot positioniert ist und nahe Station 9 liegt, um zu testen, ob es sich um einen Korallenmound handelt. Der Backengreifer kam allerdings fast leer hoch.
Stationen 11A-11G bilden ein BG-Profil in gut 20 m Wassertiefe parallel der Schelfkante von dem Mudbelt nach Norden zum nächsten geplanten Arbeitsgebiet (Paläo-Valley-Positionen). Das Material ist ein schlickartiger, gut sortierter Feinsand, ist zum Teil sehr dunkel und hat

einen starken $H_2S$-Geruch. Es ist reich an Würmern und Wurmröhren, jedoch arm an großen Karbonatschalern. Wieder kamen Fische in den Kernen mit an Deck. Der Fischreichtum der Region ist auch durch extrem hohe Anteile an Otolithen, Schuppen und Knochen manifestiert.

2.1.2007
In der Nacht kam ein starker Wind (Harmatan) mit Windstärke 8 auf und verzögerte die Arbeiten. So mußten die für die Nacht geplanten CTD-Stationen 11H-K verschoben werden. Nach Nachlassen des Windes wurden entlang eines Profils in flachen Gewässern (<50 m) bei Cap Blanc im Norden der Banc d'Arguin GKG-Kerne genommen. Im Norden der Banc d'Arguin strömt das kalte Wasser von Kanarenstrom und Upwelling auf die Banc d'Arguin. Die gewonnenen Kerne bestehen aus sauberen, hellen karbonatischen Schillen, die reich an Bivalven sind. Zwei Versuche, Vibrolot-Kerne zu nehmen, scheiterten.

3.1.2007
In der Nacht vom 2. auf den 3. Januar wurde ein CTD-Profil senkrecht zum Hang im Norden der Banc d'Arguin gemessen und Wasserproben genommen. An den CTD-Stationen wurden am Tag GKG- und SL/VL-Kerne gezogen.

Der Saharastaub hat bereits das ganze Schiff rötlich eingezuckert. Die Stimmung an Bord ist sehr gut, und die Arbeiten gehen zügig voran.

Dr. Hildegard Westphal
Fahrtleiterin auf FS Poseidon

Poseidon 346 MACUMA
2. Wochenbericht
4.-10.1.2007

In der zweiten Woche von Pos 346 wurden die Arbeiten im Bereich der Banc d'Arguin fortgesetzt. In der zweiten Hälfte der Woche verlagerten wir unser Arbeitsgebiet weiter nach Süden und untersuchten die Tiefwasser-Korallenmounds westlich Nouakchott.

4.1.2007
In der Nacht vom 3. auf den 4. Januar wurde das Nord-Süd-Profil entlang der 30 m-Isobathe der Banc d'Arguin mit 10 Backengreifer-Stationen vervollständigt, die im Norden auch in flachere Wässer (Eingang zu Baie du Levrier) reichten. Ein Trend zu einem höheren Anteil and grobkörnigem Karbonat nach Norden ist deutlich. Die Stationen am Eingang der Baie du Levrier zeigen zum Teil Grobschill, zum Teil gut sortierte Sande mit hohem Quarz-Anteil. Weiter nach Süden zeigt das Sediment eine dunkle Färbung und ist sehr feinkörnig.
Während des Tages wurden sechs Stationen auf der mittleren Banc d'Arguin beprobt, die im Bereich eines vermuteten Paläo-Valleys liegen. Aufgrund der Windverhältnisse konnte das Vibrolot erst am späten Nachmittag eingesetzt werden, so daß alle Stationen über Tag zuerst mit Backengreifer bzw. Kastengreifer beprobt wurden. Am späten Nachmittag konnten an drei der Stationen ein Schwerelot- und zwei Vibrolotkerne gezogen werden.

5.1.2007
In der Nacht vom 4. auf den 5. Januar wurden im mittleren Bereich der Banc d'Arguin ein CTD-Profil senkrecht zum Kontinentalhang gefahren. Wieder zeigte sich der Kanarenstrom deutlich in der Salinitäts- und Temperaturkurve.
Am Morgen des 5. Januar wurde das 30m-Isobathen-Profil mit Backengreifer-Stationen verdichtet. Anschließend wurden drei weitere Vibrolot-Kerne im Paläo-Valley gezogen. Danach verließen wir das Arbeitsgebiet Banc d'Arguin.

6.1.2007
Gegen Mittag erreichten wir die Banda-Mounds westlich Nouakchott. Zuerst wurden Echolotprofil-Raster gefahren, um die Korallenmounds exakt zu lokalisieren, die zum Teil weniger als 200 m breit sind. Das Echolot zeigte die Mounds sehr klar nicht nur durch ihre auffallende Morphologie, sondern auch durch die unscharfe Reflexion, die auf weichen Untergrund hinweist.

7.1.2007
In der Nacht vom 6. auf den 7. Januar wurden auf der Basis der Echolotprofile sechs CTD-Stationen gefahren. In dieser Nacht wurde unser Schiff von einer rund 50-köpfigen Delphinherde begleitet.
Am 7. Januar wurden fünf Großkastengreifer-Schwerelot-Stationen gefahren, die auf den Gipfeln der Korallenmound-Kette der Banda-Mounds positioniert waren. Die Kerne waren spektakulär in der Menge an (zum Teil lebenden) *Lophelia* und anderen Korallen. Die Muschel *Acesta*, die bislang nur aus weiter nördlich gelegenen Gebieten bekannt war, kam ebenfalls vor.

8.1.2007
In der Nacht vom 7. auf den 8. Januar wurden westlich der Banda-Mounds acht Backengreifer-Stationen gefahren. Die Stationen waren in einer Linie rund 150 m

Wassertiefe hangparallel und anschließend senkrecht den Kontinentalhang abwärts auf die Banda-Mounds zu positioniert.
Am 8. Januar wurden fünf weitere Großkastengreifer-Schwerelot-Stationen entlang der Moundkette auf den Gipfeln der Mounds gefahren. Die Kerne waren fortgesetzt spektakulär, insbesondere auch durch massenhaftes Auftreten von *Acesta*. Allerdings wurden keine weiteren lebenden Exemplare von *Lophelia* gefunden. An diesem Tag sahen wir zum ersten Mal den Horizont, der uns bislang von Sahara-Staub verhangen wurde.

9.1.2007
In der Nacht vom 8. auf den 9. Januar fuhren wir von den Banda Mounds zu den nördlichsten der Mounds in der Kette vor Nouakchott. Dort wurden Echolote gefahren, um die Korallenmounds zu lokalisieren. Danach wurden CTD-Stationen im Bereich dieser Mounds gefahren.
Am Morgen des 9.1. wurden die Korallenmounds mittels Großkastengreifer und Schwerelot beprobt. Während des anschließenden Transit zurück zur Banc d'Arguin wurden drei CTD-Profile gemessen, zwei davon in einem Gully, der den Kontinentalrand durchschneidet, eine in 40 m Wassertiefe am Rand des Südendes der Banc d'Arguin.
Anschließend wurde bei extrem ruhiger See und ruhigem Wetter ein mit vier Leuten besetztes Schlauchboot ausgesetzt, um Sedimentproben in dem Flachstwasserbereich östlich des Mudbelt der südlichen Banc d'Arguin zu nehmen. Das Schlauchboot legte rund eine Meile zurück, um in diese flachsten Bereiche zu gelangen. Die Strömung war mit geschätzten 1.5 m/s recht stark. Beprobung mittels einer Eimerdredge förderte sauberen groben Karbonatsand zutage.

10.1.2007
In der Nacht vom 9. auf den 10. Januar wurden nordöstlich des Mudbelt in rund 15 m Wassertiefe weitere Backgengreiferproben genommen, um den Datensatz zu ergänzen. Anschließend fuhren wir zur Baie du Levrier, wo ab dem Morgen Backengreifer- und Vibrolot-Stationen in Wassertiefen um 10 m gefahren wurden. Das Sediment ist überwiegend ein ausgewaschener Karbonatsand mit viel Schill, was die hohen Strömungsgeschwindigkeiten in der Baie du Levrier widerspiegelt. Bei klarer Sicht arbeiteten wir den ganzen Tag mit Blick auf Cap Blanc.

Die Arbeiten gehen weiterhin zügig voran, und die Resultate sind durchweg erfreulich.

Dr. Hildegard Westphal
Fahrtleiterin auf FS Poseidon

Poseidon 346 MACUMA
3. Wochenbericht
11.-15.1.2007

11.1.2007
In den verbleibenden zwei Arbeitstagen sollten letzte Arbeiten auf der südlichen Banc d'Arguin und dem vorgelagerten Schelfhang durchgeführt werden. In der Nacht vom 10. auf den 11. Januar wurde ein CTD-Transekt vor Cap Blanc in Ost-West-Richtung mit Wassertiefen von 50-500 m gemessen, um den Einstrom von Auftriebs- und Kanarenstrom-Wassermassen auf die Banc d'Arguin zu erfassen. Am Morgen des 11. Januar sollten die drei flachsten dieser Stationen (50-10 m Wassertiefe) mittels Vibrolot beprobt werden. Nach der ersten dieser Stationen brieste es stark auf; um 10:30 h wurden wegen Windstärke 8 die Arbeiten unterbrochen. Um 11:00 hatte der Wind bis auf Stärken von 9-10 zugenommen, so daß alle Arbeiten abgebrochen werden mußten und der Rücktransit nach Las Palmas begonnen wurde.

11.-14.1.2007
Transit Arbeitsgebiet-Las Palmas, zum Teil bei starkem Wind.

14.1.2007
Einlaufen um 9:00 und damit 24 h früher als geplant in Las Palmas.

Trotz des Abbruchs der Arbeiten verliefen alle drei Unterprojekte der Fahrt POS 346 (atypische Flachwasserkarbonate, quartäre Klimaentwicklung, Tiefwasserriffe) äußerst erfolgreich, was nicht nur die Menge an Daten- und Probenmaterial betrifft. Die Zusammenarbeit zwischen Mannschaft und Wissenschaftlern war sehr gut, die Stimmung ebenso. Unser herzlicher Dank geht an Kapitän Michael Schneider, ohne den diese Fahrt niemals so erfolgreich hätte werden können, an den Ersten Horst Hagen, an Bootsmann Achim Mischker und an die gesamte Mannschaft.

Dr. Hildegard Westphal
Fahrtleiterin auf FS Poseidon

Publications of this series:

**No. 1** **Wefer, G., E. Suess and cruise participants**
Bericht über die POLARSTERN-Fahrt ANT IV/2, Rio de Janeiro - Punta Arenas, 6.11. - 1.12.1985.
60 pages, Bremen, 1986.

**No. 2** **Hoffmann, G.**
Holozänstratigraphie und Küstenlinienverlagerung an der andalusischen Mittelmeerküste.
173 pages, Bremen, 1988. (out of print)

**No. 3** **Wefer, G. and cruise participants**
Bericht über die METEOR-Fahrt M 6/6, Libreville - Las Palmas, 18.2. - 23.3.1988.
97 pages, Bremen, 1988.

**No. 4** **Wefer, G., G.F. Lutze, T.J. Müller, O. Pfannkuche, W. Schenke, G. Siedler, W. Zenk**
Kurzbericht über die METEOR-Expedition No. 6, Hamburg - Hamburg, 28.10.1987 - 19.5.1988.
29 pages, Bremen, 1988. (out of print)

**No. 5** **Fischer, G.**
Stabile Kohlenstoff-Isotope in partikulärer organischer Substanz aus dem Südpolarmeer (Atlantischer Sektor). 161 pages, Bremen, 1989.

**No. 6** **Berger, W.H. and G. Wefer**
Partikelfluß und Kohlenstoffkreislauf im Ozean.
Bericht und Kurzfassungen über den Workshop vom 3.-4. Juli 1989 in Bremen.
57 pages, Bremen, 1989.

**No. 7** **Wefer, G. and cruise participants**
Bericht über die METEOR - Fahrt M 9/4, Dakar - Santa Cruz, 19.2. - 16.3.1989.
103 pages, Bremen, 1989.

**No. 8** **Kölling, M.**
Modellierung geochemischer Prozesse im Sickerwasser und Grundwasser.
135 pages, Bremen, 1990.

**No. 9** **Heinze, P.-M.**
Das Auftriebsgeschehen vor Peru im Spätquartär. 204 pages, Bremen, 1990. (out of print)

**No. 10** **Willems, H., G. Wefer, M. Rinski, B. Donner, H.-J. Bellmann, L. Eißmann, A. Müller, B.W. Flemming, H.-C. Höfle, J. Merkt, H. Streif, G. Hertweck, H. Kuntze, J. Schwaar, W. Schäfer, M.-G. Schulz, F. Grube, B. Menke**
Beiträge zur Geologie und Paläontologie Norddeutschlands: Exkursionsführer.
202 pages, Bremen, 1990.

**No. 11** **Wefer, G. and cruise participants**
Bericht über die METEOR-Fahrt M 12/1, Kapstadt - Funchal, 13.3.1990 - 14.4.1990.
66 pages, Bremen, 1990.

**No. 12** **Dahmke, A., H.D. Schulz, A. Kölling, F. Kracht, A. Lücke**
Schwermetallspuren und geochemische Gleichgewichte zwischen Porenlösung und Sediment im Wesermündungsgebiet. BMFT-Projekt MFU 0562, Abschlußbericht. 121 pages, Bremen, 1991.

**No. 13** **Rostek, F.**
Physikalische Strukturen von Tiefseesedimenten des Südatlantiks und ihre Erfassung in Echolotregistrierungen. 209 pages, Bremen, 1991.

**No. 14** **Baumann, M.**
Die Ablagerung von Tschernobyl-Radiocäsium in der Norwegischen See und in der Nordsee.
133 pages, Bremen, 1991. (out of print)

**No. 15** **Kölling, A.**
Frühdiagenetische Prozesse und Stoff-Flüsse in marinen und ästuarinen Sedimenten.
140 pages, Bremen, 1991.

**No. 16** **SFB 261 (ed.)**
1. Kolloquium des Sonderforschungsbereichs 261 der Universität Bremen (14.Juni 1991):
Der Südatlantik im Spätquartär: Rekonstruktion von Stoffhaushalt und Stromsystemen.
Kurzfassungen der Vorträge und Poster. 66 pages, Bremen, 1991.

**No. 17** **Pätzold, J. and cruise participants**
Bericht und erste Ergebnisse über die METEOR-Fahrt M 15/2, Rio de Janeiro - Vitoria,
18.1. - 7.2.1991. 46 pages, Bremen, 1993.

**No. 18** **Wefer, G. and cruise participants**
Bericht und erste Ergebnisse über die METEOR-Fahrt M 16/1, Pointe Noire - Recife,
27.3. - 25.4.1991. 120 pages, Bremen, 1991.

**No. 19** **Schulz, H.D. and cruise participants**
Bericht und erste Ergebnisse über die METEOR-Fahrt M 16/2, Recife - Belem, 28.4. - 20.5.1991.
149 pages, Bremen, 1991.

**No. 20** **Berner, H.**
Mechanismen der Sedimentbildung in der Fram-Straße, im Arktischen Ozean und in der Norwegischen See. 167 pages, Bremen, 1991.

**No. 21** **Schneider, R.**
Spätquartäre Produktivitätsänderungen im östlichen Angola-Becken: Reaktion auf Variationen im Passat-Monsun-Windsystem und in der Advektion des Benguela-Küstenstroms.
198 pages, Bremen, 1991. (out of print)

**No. 22** **Hebbeln, D.**
Spätquartäre Stratigraphie und Paläozeanographie in der Fram-Straße. 174 pages, Bremen, 1991.

**No. 23** **Lücke, A.**
Umsetzungsprozesse organischer Substanz während der Frühdiagenese in ästuarinen Sedimenten.
137 pages, Bremen, 1991.

**No. 24** **Wefer, G. and cruise participants**
Bericht und erste Ergebnisse der METEOR-Fahrt M 20/1, Bremen - Abidjan, 18.11.- 22.12.1991.
74 pages, Bremen, 1992.

**No. 25** **Schulz, H.D. and cruise participants**
Bericht und erste Ergebnisse der METEOR-Fahrt M 20/2, Abidjan - Dakar, 27.12.1991 - 3.2.1992.
173 pages, Bremen, 1992.

**No. 26** **Gingele, F.**
Zur klimaabhängigen Bildung biogener und terrigener Sedimente und ihrer Veränderung durch die Frühdiagenese im zentralen und östlichen Südatlantik. 202 pages, Bremen, 1992.

**No. 27** **Bickert, T.**
Rekonstruktion der spätquartären Bodenwasserzirkulation im östlichen Südatlantik über stabile Isotope benthischer Foraminiferen. 205 pages, Bremen, 1992. (out of print)

**No. 28** **Schmidt, H.**
Der Benguela-Strom im Bereich des Walfisch-Rückens im Spätquartär. 172 pages, Bremen, 1992.

**No. 29** **Meinecke, G.**
Spätquartäre Oberflächenwassertemperaturen im östlichen äquatorialen Atlantik.
181 pages, Bremen, 1992.

**No. 30** **Bathmann, U., U. Bleil, A. Dahmke, P. Müller, A. Nehrkorn, E.-M. Nöthig, M. Olesch, J. Pätzold, H.D. Schulz, V. Smetacek, V. Spieß, G. Wefer, H. Willems**
Bericht des Graduierten Kollegs. Stoff-Flüsse in marinen Geosystemen.
Berichtszeitraum Oktober 1990 - Dezember 1992. 396 pages, Bremen, 1992.

**No. 31** **Damm, E.**
Frühdiagenetische Verteilung von Schwermetallen in Schlicksedimenten der westlichen Ostsee.
115 pages, Bremen, 1992.

**No. 32** **Antia, E.E.**
Sedimentology, Morphodynamics and Facies Association of a mesotidal Barrier Island Shoreface (Spiekeroog, Southern North Sea). 370 pages, Bremen, 1993.

**No. 33** **Duinker, J. and G. Wefer (ed.)**
Bericht über den 1. JGOFS-Workshop. 1./2. Dezember 1992 in Bremen. 83 pages, Bremen, 1993.

**No. 34** **Kasten, S.**
Die Verteilung von Schwermetallen in den Sedimenten eines stadtbremischen Hafenbeckens.
103 pages, Bremen, 1993.

**No. 35** **Spieß, V.**
Digitale Sedimentographie. Neue Wege zu einer hochauflösenden Akustostratigraphie.
199 pages, Bremen, 1993.

**No. 36** **Schinzel, U.**
Laborversuche zu frühdiagenetischen Reaktionen von Eisen (III) - Oxidhydraten in marinen Sedimenten.189 pages, Bremen, 1993.

**No. 37** **Sieger, R.**
CoTAM - ein Modell zur Modellierung des Schwermetalltransports in Grundwasserleitern.
56 pages, Bremen, 1993. (out of print)

**No. 38** **Willems, H. (ed.)**
Geoscientific Investigations in the Tethyan Himalayas. 183 pages, Bremen, 1993.

**No. 39** **Hamer, K.**
Entwicklung von Laborversuchen als Grundlage für die Modellierung des Transportverhaltens von Arsenat, Blei, Cadmium und Kupfer in wassergesättigten Säulen. 147 pages, Bremen, 1993.

**No. 40** **Sieger, R.**
Modellierung des Stofftransports in porösen Medien unter Ankopplung kinetisch gesteuerter Sorptions- und Redoxprozesse sowie thermischer Gleichgewichte. 158 pages, Bremen, 1993.

**No. 41** **Thießen, W.**
Magnetische Eigenschaften von Sedimenten des östlichen Südatlantiks und ihre paläozeanographische Relevanz. 170 pages, Bremen, 1993.

**No. 42** **Spieß, V. and cruise participants**
Report and preliminary results of METEOR-Cruise M 23/1, Kapstadt - Rio de Janeiro, 4.-25.2.1993. 139 pages, Bremen, 1994.

**No. 43** **Bleil, U. and cruise participants**
Report and preliminary results of METEOR-Cruise M 23/2, Rio de Janeiro - Recife, 27.2.-19.3.1993 133 pages, Bremen, 1994.

**No. 44** **Wefer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 23/3, Recife - Las Palmas, 21.3. - 12.4.1993 71 pages, Bremen, 1994.

**No. 45** **Giese, M. and G. Wefer (ed.)**
Bericht über den 2. JGOFS-Workshop. 18../19. November 1993 in Bremen. 93 pages, Bremen, 1994.

**No. 46** **Balzer, W. and cruise participants**
Report and preliminary results of METEOR-Cruise M 22/1, Hamburg - Recife, 22.9. - 21.10.1992. 24 pages, Bremen, 1994.

**No. 47** **Stax, R.**
Zyklische Sedimentation von organischem Kohlenstoff in der Japan See: Anzeiger für Änderungen von Paläoozeanographie und Paläoklima im Spätkänozoikum. 150 pages, Bremen, 1994.

**No. 48** **Skowronek, F.**
Frühdiagenetische Stoff-Flüsse gelöster Schwermetalle an der Oberfläche von Sedimenten des Weser Ästuares. 107 pages, Bremen, 1994.

**No. 49** **Dersch-Hansmann, M.**
Zur Klimaentwicklung in Ostasien während der letzten 5 Millionen Jahre: Terrigener Sedimenteintrag in die Japan See (ODP Ausfahrt 128). 149 pages, Bremen, 1994.

**No. 50** **Zabel, M.**
Frühdiagenetische Stoff-Flüsse in Oberflächen-Sedimenten des äquatorialen und östlichen Südatlantik. 129 pages, Bremen, 1994.

**No. 51** **Bleil, U. and cruise participants**
Report and preliminary results of SONNE-Cruise SO 86, Buenos Aires - Capetown, 22.4. - 31.5.93 116 pages, Bremen, 1994.

**No. 52** **Symposium: The South Atlantic: Present and Past Circulation.**
Bremen, Germany, 15 - 19 August 1994. Abstracts. 167 pages, Bremen, 1994.

**No. 53** **Kretzmann, U.B.**
$^{57}$Fe-Mössbauer-Spektroskopie an Sedimenten - Möglichkeiten und Grenzen. 183 pages, Bremen, 1994.

**No. 54** **Bachmann, M.**
Die Karbonatrampe von Organyà im oberen Oberapt und unteren Unteralb (NE-Spanien, Prov. Lerida): Fazies, Zyklo- und Sequenzstratigraphie. 147 pages, Bremen, 1994. (out of print)

**No. 55** **Kemle-von Mücke, S.**
Oberflächenwasserstruktur und -zirkulation des Südostatlantiks im Spätquartär. 151 pages, Bremen, 1994.

**No. 56** **Petermann, H.**
Magnetotaktische Bakterien und ihre Magnetosome in Oberflächensedimenten des Südatlantiks. 134 pages, Bremen, 1994.

**No. 57** **Mulitza, S.**
Spätquartäre Variationen der oberflächennahen Hydrographie im westlichen äquatorialen Atlantik. 97 pages, Bremen, 1994.

**No. 58** **Segl, M. and cruise participants**
Report and preliminary results of METEOR-Cruise M 29/1, Buenos-Aires - Montevideo, 17.6. - 13.7.1994
94 pages, Bremen, 1994.

**No. 59** **Bleil, U. and cruise participants**
Report and preliminary results of METEOR-Cruise M 29/2, Montevideo - Rio de Janiero 15.7. - 8.8.1994. 153 pages, Bremen, 1994.

**No. 60** **Henrich, R. and cruise participants**
Report and preliminary results of METEOR-Cruise M 29/3, Rio de Janeiro - Las Palmas 11.8. - 5.9.1994. Bremen, 1994. (out of print)

**No. 61** **Sagemann, J.**
Saisonale Variationen von Porenwasserprofilen, Nährstoff-Flüssen und Reaktionen in intertidalen Sedimenten des Weser-Ästuars. 110 pages, Bremen, 1994. (out of print)

**No. 62** **Giese, M. and G. Wefer**
Bericht über den 3. JGOFS-Workshop. 5./6. Dezember 1994 in Bremen.
84 pages, Bremen, 1995.

**No. 63** **Mann, U.**
Genese kretazischer Schwarzschiefer in Kolumbien: Globale vs. regionale/lokale Prozesse.
153 pages, Bremen, 1995. (out of print)

**No. 64** **Willems, H., Wan X., Yin J., Dongdui L., Liu G., S. Dürr, K.-U. Gräfe**
The Mesozoic development of the N-Indian passive margin and of the Xigaze Forearc Basin in southern Tibet, China. – Excursion Guide to IGCP 362 Working-Group Meeting "Integrated Stratigraphy". 113 pages, Bremen, 1995. (out of print)

**No. 65** **Hünken, U.**
Liefergebiets - Charakterisierung proterozoischer Goldseifen in Ghana anhand von Fluideinschluß - Untersuchungen. 270 pages, Bremen, 1995.

**No. 66** **Nyandwi, N.**
The Nature of the Sediment Distribution Patterns in ther Spiekeroog Backbarrier Area, the East Frisian Islands. 162 pages, Bremen, 1995.

**No. 67** **Isenbeck-Schröter, M.**
Transportverhalten von Schwermetallkationen und Oxoanionen in wassergesättigten Sanden. - Laborversuche in Säulen und ihre Modellierung -. 182 pages, Bremen, 1995.

**No. 68** **Hebbeln, D. and cruise participants**
Report and preliminary results of SONNE-Cruise SO 102, Valparaiso - Valparaiso, 95.
134 pages, Bremen, 1995.

**No. 69** **Willems, H. (Sprecher), U.Bathmann, U. Bleil, T. v. Dobeneck, K. Herterich, B.B. Jorgensen, E.-M. Nöthig, M. Olesch, J. Pätzold, H.D. Schulz, V. Smetacek, V. Speiß. G. Wefer**
Bericht des Graduierten-Kollegs Stoff-Flüsse in marine Geosystemen.
Berichtszeitraum Januar 1993 - Dezember 1995.
45 & 468 pages, Bremen, 1995.

**No. 70** **Giese, M. and G. Wefer**
Bericht über den 4. JGOFS-Workshop. 20./21. November 1995 in Bremen. 60 pages, Bremen, 1996. (out of print)

**No. 71** **Meggers, H.**
Pliozän-quartäre Karbonatsedimentation und Paläozeanographie des Nordatlantiks und des Europäischen Nordmeeres - Hinweise aus planktischen Foraminiferengemeinschaften.
143 pages, Bremen, 1996. (out of print)

**No. 72** **Teske, A.**
Phylogenetische und ökologische Untersuchungen an Bakterien des oxidativen und reduktiven marinen Schwefelkreislaufs mittels ribosomaler RNA. 220 pages, Bremen, 1996. (out of print)

**No. 73** **Andersen, N.**
Biogeochemische Charakterisierung von Sinkstoffen und Sedimenten aus ostatlantischen Produktions-Systemen mit Hilfe von Biomarkern. 215 pages, Bremen, 1996.

**No. 74** **Treppke, U.**
Saisonalität im Diatomeen- und Silikoflagellatenfluß im östlichen tropischen und subtropischen Atlantik. 200 pages, Bremen, 1996.

**No. 75** **Schüring, J.**
Die Verwendung von Steinkohlebergematerialien im Deponiebau im Hinblick auf die Pyritverwitterung und die Eignung als geochemische Barriere. 110 pages, Bremen, 1996.

**No. 76** **Pätzold, J. and cruise participants**
Report and preliminary results of VICTOR HENSEN cruise JOPS II, Leg 6, Fortaleza - Recife, 10.3. - 26.3. 1995 and Leg 8, Vítoria - Vítoria, 10.4. - 23.4.1995.
87 pages, Bremen, 1996.

**No. 77** **Bleil, U. and cruise participants**
Report and preliminary results of METEOR-Cruise M 34/1, Cape Town - Walvis Bay, 3.-26.1.1996.
129 pages, Bremen, 1996.

**No. 78** **Schulz, H.D. and cruise participants**
Report and preliminary results of METEOR-Cruise M 34/2, Walvis Bay - Walvis Bay, 29.1.-18.2.96
133 pages, Bremen, 1996.

**No. 79** **Wefer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 34/3, Walvis Bay - Recife, 21.2.-17.3.1996.
168 pages, Bremen, 1996.

**No. 80** **Fischer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 34/4, Recife - Bridgetown, 19.3.-15.4.1996.
105 pages, Bremen, 1996.

**No. 81** **Kulbrok, F.**
Biostratigraphie, Fazies und Sequenzstratigraphie einer Karbonatrampe in den Schichten der Oberkreide und des Alttertiärs Nordost-Ägyptens (Eastern Desert, N'Golf von Suez, Sinai).
153 pages, Bremen, 1996.

**No. 82** **Kasten, S.**
Early Diagenetic Metal Enrichments in Marine Sediments as Documents of Nonsteady-State Depositional Conditions. Bremen, 1996.

**No. 83** **Holmes, M.E.**
Reconstruction of Surface Ocean Nitrate Utilization in the Southeast Atlantic Ocean Based on Stable Nitrogen Isotopes. 113 pages, Bremen, 1996.

**No. 84** **Rühlemann, C.**
Akkumulation von Carbonat und organischem Kohlenstoff im tropischen Atlantik: Spätquartäre Produktivitäts-Variationen und ihre Steuerungsmechanismen.
139 pages, Bremen, 1996.

**No. 85** **Ratmeyer, V.**
Untersuchungen zum Eintrag und Transport lithogener und organischer partikulärer Substanz im östlichen subtropischen Nordatlantik. 154 pages, Bremen, 1996.

**No. 86** **Cepek, M.**
Zeitliche und räumliche Variationen von Coccolithophoriden-Gemeinschaften im subtropischen Ost-Atlantik: Untersuchungen an Plankton, Sinkstoffen und Sedimenten.
156 pages, Bremen, 1996.

**No. 87** **Otto, S.**
Die Bedeutung von gelöstem organischen Kohlenstoff (DOC) für den Kohlenstoffffluß im Ozean.
150 pages, Bremen, 1996.

**No. 88** **Hensen, C.**
Frühdiagenetische Prozesse und Quantifizierung benthischer Stoff-Flüsse in Oberflächensedimenten des Südatlantiks.
132 pages, Bremen, 1996.

**No. 89** **Giese, M. and G. Wefer**
Bericht über den 5. JGOFS-Workshop. 27./28. November 1996 in Bremen. 73 pages, Bremen, 1997.

**No. 90** **Wefer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 37/1, Lisbon - Las Palmas, 4.-23.12.1996.
79 pages, Bremen, 1997.

**No. 91** **Isenbeck-Schröter, M., E. Bedbur, M. Kofod, B. König, T. Schramm & G. Mattheß**
Occurrence of Pesticide Residues in Water - Assessment of the Current Situation in Selected EU Countries. 65 pages, Bremen 1997.

**No. 92** **Kühn, M.**
Geochemische Folgereaktionen bei der hydrogeothermalen Energiegewinnung.
129 pages, Bremen 1997.

**No. 93** **Determann, S. & K. Herterich**
JGOFS-A6 "Daten und Modelle": Sammlung JGOFS-relevanter Modelle in Deutschland.
26 pages, Bremen, 1997.

**No. 94** **Fischer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 38/1, Las Palmas - Recife, 25.1.-1.3.1997, with Appendix: Core Descriptions from METEOR Cruise M 37/1. Bremen, 1997.

**No. 95** **Bleil, U. and cruise participants**
Report and preliminary results of METEOR-Cruise M 38/2, Recife - Las Palmas, 4.3.-14.4.1997.
126 pages, Bremen, 1997.

**No. 96** **Neuer, S. and cruise participants**
Report and preliminary results of VICTOR HENSEN-Cruise 96/1. Bremen, 1997.

**No. 97** **Villinger, H. and cruise participants**
Fahrtbericht SO 111, 20.8. - 16.9.1996. 115 pages, Bremen, 1997.

**No. 98** **Lüning, S.**
Late Cretaceous - Early Tertiary sequence stratigraphy, paleoecology and geodynamics of Eastern Sinai, Egypt. 218 pages, Bremen, 1997.

**No. 99** **Haese, R.R.**
Beschreibung und Quantifizierung frühdiagenetischer Reaktionen des Eisens in Sedimenten des Südatlantiks. 118 pages, Bremen, 1997.

**No. 100** **Lührte, R. von**
Verwertung von Bremer Baggergut als Material zur Oberflächenabdichtung von Deponien - Geochemisches Langzeitverhalten und Schwermetall-Mobilität (Cd, Cu, Ni, Pb, Zn). Bremen, 1997.

**No. 101** **Ebert, M.**
Der Einfluß des Redoxmilieus auf die Mobilität von Chrom im durchströmten Aquifer. 135 pages, Bremen, 1997.

**No. 102** **Krögel, F.**
Einfluß von Viskosität und Dichte des Seewassers auf Transport und Ablagerung von Wattsedimenten (Langeooger Rückseitenwatt, südliche Nordsee). 168 pages, Bremen, 1997.

**No. 103** **Kerntopf, B.**
Dinoflagellate Distribution Patterns and Preservation in the Equatorial Atlantic and Offshore North-West Africa. 137 pages, Bremen, 1997.

**No. 104** **Breitzke, M.**
Elastische Wellenausbreitung in marinen Sedimenten - Neue Entwicklungen der Ultraschall Sedimentphysik und Sedimentechographie. 298 pages, Bremen, 1997.

**No. 105** **Marchant, M.**
Rezente und spätquartäre Sedimentation planktischer Foraminiferen im Peru-Chile Strom. 115 pages, Bremen, 1997.

**No. 106** **Habicht, K.S.**
Sulfur isotope fractionation in marine sediments and bacterial cultures. 125 pages, Bremen, 1997.

**No. 107** **Hamer, K., R.v. Lührte, G. Becker, T. Felis, S. Keffel, B. Strotmann, C. Waschkowitz, M. Kölling, M. Isenbeck-Schröter, H.D. Schulz**
Endbericht zum Forschungsvorhaben 060 des Landes Bremen: Baggergut der Hafengruppe Bremen-Stadt: Modelluntersuchungen zur Schwermetallmobilität und Möglichkeiten der Verwertung von Hafenschlick aus Bremischen Häfen. 98 pages, Bremen, 1997.

**No. 108** **Greeff, O.W.**
Entwicklung und Erprobung eines benthischen Landersystemes zur *in situ*-Bestimmung von Sulfatreduktionsraten mariner Sedimente. 121 pages, Bremen, 1997.

**No. 109** **Pätzold, M. und G. Wefer**
Bericht über den 6. JGOFS-Workshop am 4./5.12.1997 in Bremen. Im Anhang: Publikationen zum deutschen Beitrag zur Joint Global Ocean Flux Study (JGOFS), Stand 1/1998. 122 pages, Bremen, 1998.

**No. 110** **Landenberger, H.**
CoTReM, ein Multi-Komponenten Transport- und Reaktions-Modell. 142 pages, Bremen, 1998.

**No. 111** **Villinger, H. und Fahrtteilnehmer**
Fahrtbericht SO 124, 4.10. - 16.10.199. 90 pages, Bremen, 1997.

**No. 112** **Gietl, R.**
Biostratigraphie und Sedimentationsmuster einer nordostägyptischen Karbonatrampe unter Berücksichtigung der Alveolinen-Faunen. 142 pages, Bremen, 1998.

**No. 113** **Ziebis, W.**
The Impact of the Thalassinidean Shrimp *Callianassa truncata* on the Geochemistry of permeable, coastal Sediments. 158 pages, Bremen 1998.

**No. 114** **Schulz, H.D. and cruise participants**
Report and preliminary results of METEOR-Cruise M 41/1, Málaga - Libreville, 13.2.-15.3.1998. Bremen, 1998.

**No. 115** **Völker, D.J.**
Untersuchungen an strömungsbeeinflußten Sedimentationsmustern im Südozean. Interpretation sedimentechographischer Daten und numerische Modellierung. 152 pages, Bremen, 1998.

**No. 116** **Schlünz, B.**
Riverine Organic Carbon Input into the Ocean in Relation to Late Quaternary Climate Change. 136 pages, Bremen, 1998.

**No. 117** **Kuhnert, H.**
Aufzeichnug des Klimas vor Westaustralien in stabilen Isotopen in Korallenskeletten. 109 pages, Bremen, 1998.

**No. 118** **Kirst, G.**
Rekonstruktion von Oberflächenwassertemperaturen im östlichen Südatlantik anhand von Alkenonen. 130 pages, Bremen, 1998.

**No. 119** **Dürkoop, A.**
Der Brasil-Strom im Spätquartär: Rekonstruktion der oberflächennahen Hydrographie während der letzten 400 000 Jahre. 121 pages, Bremen, 1998.

**No. 120** **Lamy, F.**
Spätquartäre Variationen des terrigenen Sedimenteintrags entlang des chilenischen Kontinentalhangs als Abbild von Klimavariabilität im Milankovi_- und Sub-Milankovi_-Zeitbereich. 141 pages, Bremen, 1998.

**No. 121** **Neuer, S. and cruise participants**
Report and preliminary results of POSEIDON-Cruise Pos 237/2, Vigo – Las Palmas, 18.3.-31.3.1998. 39 pages, Bremen, 1998

**No. 122** **Romero, O.E.**
Marine planktonic diatoms from the tropical and equatorial Atlantic: temporal flux patterns and the sediment record. 205 pages, Bremen, 1998.

**No. 123** **Spiess, V. und Fahrtteilnehmer**
Report and preliminary results of RV SONNE Cruise 125, Cochin – Chittagong, 17.10.-17.11.1997. 128 pages, Bremen, 1998.

**No. 124** **Arz, H.W.**
Dokumentation von kurzfristigen Klimaschwankungen des Spätquartärs in Sedimenten des westlichen äquatorialen Atlantiks. 96 pages, Bremen, 1998.

**No. 125** **Wolff, T.**
Mixed layer characteristics in the equatorial Atlantic during the late Quaternary as deduced from planktonic foraminifera. 132 pages, Bremen, 1998.

**No. 126** **Dittert, N.**
Late Quaternary Planktic Foraminifera Assemblages in the South Atlantic Ocean: Quantitative Determination and Preservational Aspects. 165 pages, Bremen, 1998.

**No. 127** **Höll, C.**
Kalkige und organisch-wandige Dinoflagellaten-Zysten in Spätquartären Sedimenten des tropischen Atlantiks und ihre palökologische Auswertbarkeit. 121 pages, Bremen, 1998.

**No. 128** **Hencke, J.**
Redoxreaktionen im Grundwasser: Etablierung und Verlagerung von Reaktionsfronten und ihre Bedeutung für die Spurenelement-Mobilität. 122 pages, Bremen 1998.

**No. 129** **Pätzold, J. and cruise participants**
Report and preliminary results of METEOR-Cruise M 41/3, Vítoria, Brasil – Salvador de Bahia, Brasil, 18.4. - 15.5.1998. Bremen, 1999.

**No. 130** **Fischer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 41/4, Salvador de Bahia, Brasil – Las Palmas, Spain, 18.5. – 13.6.1998. Bremen, 1999.

**No. 131** **Schlünz, B. und G. Wefer**
Bericht über den 7. JGOFS-Workshop am 3. und 4.12.1998 in Bremen. Im Anhang: Publikationen zum deutschen Beitrag zur Joint Global Ocean Flux Study (JGOFS), Stand 1/ 1999. 100 pages, Bremen, 1999.

**No. 132** **Wefer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 42/4, Las Palmas - Las Palmas - Viena do Castelo; 26.09.1998 - 26.10.1998. 104 pages, Bremen, 1999.

**No. 133** **Felis, T.**
Climate and ocean variability reconstructed from stable isotope records of modern subtropical corals (Northern Red Sea). 111 pages, Bremen, 1999.

**No. 134** **Draschba , S.**
North Atlantic climate variability recorded in reef corals from Bermuda. 108 pages, Bremen, 1999.

**No. 135** **Schmieder, F.**
Magnetic Cyclostratigraphy of South Atlantic Sediments. 82 pages, Bremen, 1999.

**No. 136** **Rieß, W.**
In situ measurements of respiration and mineralisation processes – Interaction between fauna and geochemical fluxes at active interfaces. 68 pages, Bremen, 1999.

**No. 137** **Devey, C.W. and cruise participants**
Report and shipboard results from METEOR-cruise M 41/2, Libreville – Vitoria, 18.3. – 15.4.98. 59 pages, Bremen, 1999.

**No. 138** **Wenzhöfer, F.**
Biogeochemical processes at the sediment water interface and quantification of metabolically driven calcite dissolution in deep sea sediments. 103 pages, Bremen, 1999.

**No. 139** **Klump, J.**
Biogenic barite as a proxy of paleoproductivity variations in the Southern Peru-Chile Current. 107 pages, Bremen, 1999.

**No. 140** **Huber, R.**
Carbonate sedimentation in the northern Northatlantic since the late pliocene. 103 pages, Bremen, 1999.

**No. 141** **Schulz, H.**
Nitrate-storing sulfur bacteria in sediments of coastal upwelling. 94 pages, Bremen, 1999.

**No. 142** **Mai, S.**
Die Sedimentverteilung im Wattenmeer: ein Simulationsmodell. 114 pages, Bremen, 1999.

**No. 143** **Neuer, S. and cruise participants**
Report and preliminary results of Poseidon Cruise 248, Las Palmas - Las Palmas, 15.2.-26.2.1999.
45 pages, Bremen, 1999.

**No. 144** **Weber, A.**
Schwefelkreislauf in marinen Sedimenten und Messung von *in situ* Sulfatreduktionsraten.
122 pages, Bremen, 1999.

**No. 145** **Hadeler, A.**
Sorptionsreaktionen im Grundwasser: Unterschiedliche Aspekte bei der Modellierung des Transportverhaltens von Zink. 122 pages, 1999.

**No. 146** **Dierßen, H.**
Zum Kreislauf ausgewählter Spurenmetalle im Südatlantik: Vertikaltransport und Wechselwirkung zwischen Partikeln und Lösung. 167 pages, Bremen, 1999.

**No. 147** **Zühlsdorff, L.**
High resolution multi-frequency seismic surveys at the Eastern Juan de Fuca Ridge Flank and the Cascadia Margin – Evidence for thermally and tectonically driven fluid upflow in marine sediments. 118 pages, Bremen 1999.

**No. 148** **Kinkel, H.**
Living and late Quaternary Coccolithophores in the equatorial Atlantic Ocean: response of distribution and productivity patterns to changing surface water circulation. 183 pages, Bremen, 2000.

**No. 149** **Pätzold, J. and cruise participants**
Report and preliminary results of METEOR Cruise M 44/3, Aqaba (Jordan) - Safaga (Egypt) – Dubá (Saudi Arabia) – Suez (Egypt) - Haifa (Israel), 12.3.-26.3.-2.4.-4.4.1999. 135 pages, Bremen, 2000.

**No. 150** **Schlünz, B. and G. Wefer**
Bericht über den 8. JGOFS-Workshop am 2. und 3.12.1999 in Bremen. Im Anhang:
Publikationen zum deutschen Beitrag zur Joint Global Ocean Flux Study (JGOFS), Stand 1/ 2000.
95 pages, Bremen, 2000.

**No. 151** **Schnack, K.**
Biostratigraphie und fazielle Entwicklung in der Oberkreide und im Alttertiär im Bereich der Kharga Schwelle, Westliche Wüste, SW-Ägypten. 142 pages, Bremen, 2000.

**No. 152** **Karwath, B.**
Ecological studies on living and fossil calcareous dinoflagellates of the equatorial and tropical Atlantic Ocean. 175 pages, Bremen, 2000.

**No. 153** **Moustafa, Y.**
Paleoclimatic reconstructions of the Northern Red Sea during the Holocene inferred from stable isotope records of modern and fossil corals and molluscs. 102 pages, Bremen, 2000.

**No. 154** **Villinger, H. and cruise participants**
Report and preliminary results of SONNE-cruise 145-1 Balboa – Talcahuana, 21.12.1999 – 28.01.2000.
147 pages, Bremen, 2000.

**No. 155** **Rusch, A.**
Dynamik der Feinfraktion im Oberflächenhorizont permeabler Schelfsedimente. 102 pages, Bremen, 2000.

**No. 156** **Moos, C.**
Reconstruction of upwelling intensity and paleo-nutrient gradients in the northwest Arabian Sea derived from stable carbon and oxygen isotopes of planktic foraminifera. 103 pages, Bremen, 2000.

**No. 157** **Xu, W.**
Mass physical sediment properties and trends in a Wadden Sea tidal basin. 127 pages, Bremen, 2000.

**No. 158** **Meinecke, G. and cruise participants**
Report and preliminary results of METEOR Cruise M 45/1, Malaga (Spain) - Lissabon (Portugal),
19.05. - 08.06.1999. 39 pages, Bremen, 2000.

**No. 159** **Vink, A.**
Reconstruction of recent and late Quaternary surface water masses of the western subtropical Atlantic Ocean based on calcareous and organic-walled dinoflagellate cysts. 160 pages, Bremen, 2000.

**No. 160** **Willems, H. (Sprecher), U. Bleil, R. Henrich, K. Herterich, B.B. Jørgensen, H.-J. Kuß, M. Olesch, H.D. Schulz,V. Spieß, G. Wefer**
Abschlußbericht des Graduierten-Kollegs Stoff-Flüsse in marine Geosystemen.
Zusammenfassung und Berichtszeitraum Januar 1996 - Dezember 2000. 340 pages, Bremen, 2000.

**No. 161** **Sprengel, C.**
Untersuchungen zur Sedimentation und Ökologie von Coccolithophoriden im Bereich der Kanarischen Inseln: Saisonale Flussmuster und Karbonatexport. 165 pages, Bremen, 2000.

**No. 162** **Donner, B. and G. Wefer**
Bericht über den JGOFS-Workshop am 18.-21.9.2000 in Bremen:
Biogeochemical Cycles: German Contributions to the International Joint Global Ocean Flux Study.
87 pages, Bremen, 2000.

**No. 163** **Neuer, S. and cruise participants**
Report and preliminary results of Meteor Cruise M 45/5, Bremen – Las Palmas, October 1 – November 3, 1999. 93 pages, Bremen, 2000.

**No. 164** **Devey, C. and cruise participants**
Report and preliminary results of Sonne Cruise SO 145/2, Talcahuano (Chile) - Arica (Chile), February 4 – February 29, 2000. 63 pages, Bremen, 2000.

**No. 165** **Freudenthal, T.**
Reconstruction of productivity gradients in the Canary Islands region off Morocco by means of sinking particles and sediments. 147 pages, Bremen, 2000.

**No. 166** **Adler, M.**
Modeling of one-dimensional transport in porous media with respect to simultaneous geochemical reactions in CoTReM. 147 pages, Bremen, 2000.

**No. 167** **Santamarina Cuneo, P.**
Fluxes of suspended particulate matter through a tidal inlet of the East Frisian Wadden Sea (southern North Sea). 91 pages, Bremen, 2000.

**No. 168** **Benthien, A.**
Effects of $CO_2$ and nutrient concentration on the stable carbon isotope composition of C37:2 alkenones in sediments of the South Atlantic Ocean. 104 pages, Bremen, 2001.

**No. 169** **Lavik, G.**
Nitrogen isotopes of sinking matter and sediments in the South Atlantic. 140 pages, Bremen, 2001.

**No. 170** **Budziak, D.**
Late Quaternary monsoonal climate and related variations in paleoproductivity and alkenone-derived sea-surface temperatures in the western Arabian Sea. 114 pages, Bremen, 2001.

**No. 171** **Gerhardt, S.**
Late Quaternary water mass variability derived from the pteropod preservation state in sediments of the western South Atlantic Ocean and the Caribbean Sea. 109 pages, Bremen, 2001.

**No. 172** **Bleil, U. and cruise participants**
Report and preliminary results of Meteor Cruise M 46/3, Montevideo (Uruguay) – Mar del Plata (Argentina), January 4 – February 7, 2000. Bremen, 2001.

**No. 173** **Wefer, G. and cruise participants**
Report and preliminary results of Meteor Cruise M 46/4, Mar del Plata (Argentina) – Salvador da Bahia (Brazil), February 10 – March 13, 2000. With partial results of METEOR cruise M 46/2. 136 pages, Bremen, 2001.

**No. 174** **Schulz, H.D. and cruise participants**
Report and preliminary results of Meteor Cruise M 46/2, Recife (Brazil) – Montevideo (Uruguay), December 2 – December 29, 1999. 107 pages, Bremen, 2001.

**No. 175** **Schmidt, A.**
Magnetic mineral fluxes in the Quaternary South Atlantic: Implications for the paleoenvironment.
97 pages, Bremen, 2001.

**No. 176** **Bruhns, P.**
Crystal chemical characterization of heavy metal incorporation in brick burning processes.
93 pages, Bremen, 2001.

**No. 177** **Karius, V.**
Baggergut der Hafengruppe Bremen-Stadt in der Ziegelherstellung. 131 pages, Bremen, 2001.

**No. 178** **Adegbie, A. T.**
Reconstruction of paleoenvironmental conditions in Equatorial Atlantic and the Gulf of Guinea Basins for the last 245,000 years. 113 pages, Bremen, 2001.

**No. 179** **Spieß, V. and cruise participants**
Report and preliminary results of R/V Sonne Cruise SO 149, Victoria - Victoria, 16.8. - 16.9.2000.
100 pages, Bremen, 2001.

**No. 180** **Kim, J.-H.**
Reconstruction of past sea-surface temperatures in the eastern South Atlantic and the eastern South Pacific across Termination I based on the Alkenone Method. 114 pages, Bremen, 2001.

**No. 181** **von Lom-Keil, H.**
Sedimentary waves on the Namibian continental margin and in the Argentine Basin – Bottom flow reconstructions based on high resolution echosounder data. 126 pages, Bremen, 2001.

**No. 182** **Hebbeln, D. and cruise participants**
PUCK: Report and preliminary results of R/V Sonne Cruise SO 156, Valparaiso (Chile) - Talcahuano (Chile), March 29 - May 14, 2001. 195 pages, Bremen, 2001.

**No. 183** **Wendler, J.**
Reconstruction of astronomically-forced cyclic and abrupt paleoecological changes in the Upper Cretaceous Boreal Realm based on calcareous dinoflagellate cysts. 149 pages, Bremen, 2001.

**No. 184** **Volbers, A.**
Planktic foraminifera as paleoceanographic indicators: production, preservation, and reconstruction of upwelling intensity. Implications from late Quaternary South Atlantic sediments. 122 pages, Bremen, 2001.

**No. 185** **Bleil, U. and cruise participants**
Report and preliminary results of R/V METEOR Cruise M 49/3, Montevideo (Uruguay) - Salvador (Brasil), March 9 - April 1, 2001. 99 pages, Bremen, 2001.

**No. 186** **Scheibner, C.**
Architecture of a carbonate platform-to-basin transition on a structural high (Campanian-early Eocene, Eastern Desert, Egypt) – classical and modelling approaches combined. 173 pages, Bremen, 2001.

**No. 187** **Schneider, S.**
Quartäre Schwankungen in Strömungsintensität und Produktivität als Abbild der Wassermassen-Variabilität im äquatorialen Atlantik (ODP Sites 959 und 663): Ergebnisse aus Siltkorn-Analysen. 134 pages, Bremen, 2001.

**No. 188** **Uliana, E.**
Late Quaternary biogenic opal sedimentation in diatom assemblages in Kongo Fan sediments. 96 pages, Bremen, 2002.

**No. 189** **Esper, O.**
Reconstruction of Recent and Late Quaternary oceanographic conditions in the eastern South Atlantic Ocean based on calcareous- and organic-walled dinoflagellate cysts. 130 pages, Bremen, 2001.

**No. 190** **Wendler, I.**
Production and preservation of calcareous dinoflagellate cysts in the modern Arabian Sea. 117 pages, Bremen, 2002.

**No. 191** **Bauer, J.**
Late Cenomanian – Santonian carbonate platform evolution of Sinai (Egypt): stratigraphy, facies, and sequence architecture. 178 pages, Bremen, 2002.

**No. 192** **Hildebrand-Habel, T.**
Die Entwicklung kalkiger Dinoflagellaten im Südatlantik seit der höheren Oberkreide. 152 pages, Bremen, 2002.

**No. 193** **Hecht, H.**
Sauerstoff-Optopoden zur Quantifizierung von Pyritverwitterungsprozessen im Labor- und Langzeit-in-situ-Einsatz. Entwicklung - Anwendung – Modellierung. 130 pages, Bremen, 2002.

**No. 194** **Fischer, G. and cruise participants**
Report and Preliminary Results of RV METEOR-Cruise M49/4, Salvador da Bahia – Halifax, 4.4.-5.5.2001. 84 pages, Bremen, 2002.

**No. 195** **Gröger, M.**
Deep-water circulation in the western equatorial Atlantic: inferences from carbonate preservation studies and silt grain-size analysis. 95 pages, Bremen, 2002.

**No. 196** **Meinecke,G. and cruise participants**
Report of RV POSEIDON Cruise POS 271, Las Palmas - Las Palmas, 19.3.-29.3.2001. 19 pages, Bremen, 2002.

**No. 197** **Meggers, H. and cruise participants**
Report of RV POSEIDON Cruise POS 272, Las Palmas - Las Palmas, 1.4.-14.4.2001. 19 pages, Bremen, 2002.

**No. 198** **Gräfe, K.-U.**
Stratigraphische Korrelation und Steuerungsfaktoren Sedimentärer Zyklen in ausgewählten Borealen und Tethyalen Becken des Cenoman/Turon (Oberkreide) Europas und Nordwestafrikas. 197 pages, Bremen, 2002.

**No. 199** **Jahn, B.**
Mid to Late Pleistocene Variations of Marine Productivity in and Terrigenous Input to the Southeast Atlantic. 97 pages, Bremen, 2002.

**No. 200** **Al-Rousan, S.**
Ocean and climate history recorded in stable isotopes of coral and foraminifers from the northern Gulf of Aqaba. 116 pages, Bremen, 2002.

**No. 201** **Azouzi, B.**
Regionalisierung hydraulischer und hydrogeochemischer Daten mit geostatistischen Methoden. 108 pages, Bremen, 2002.

**No. 202** **Spieß, V. and cruise participants**
Report and preliminary results of METEOR Cruise M 47/3, Libreville (Gabun) - Walvis Bay (Namibia), 01.06 - 03.07.2000. 70 pages, Bremen 2002.

**No. 203** **Spieß, V. and cruise participants**
Report and preliminary results of METEOR Cruise M 49/2, Montevideo (Uruguay) - Montevideo, 13.02 - 07.03.2001. 84 pages, Bremen 2002.

**No. 204** **Mollenhauer, G.**
Organic carbon accumulation in the South Atlantic Ocean: Sedimentary processes and glacial/interglacial Budgets. 139 pages, Bremen 2002.

**No. 205** **Spieß, V. and cruise participants**
Report and preliminary results of METEOR Cruise M49/1, Cape Town (South Africa) - Montevideo (Uruguay), 04.01.2001 - 10.02.2001. 57 pages, Bremen, 2003.

**No. 206** **Meier, K.J.S.**
Calcareous dinoflagellates from the Mediterranean Sea: taxonomy, ecology and palaeoenvironmental application. 126 pages, Bremen, 2003.

**No. 207** **Rakic, S.**
Untersuchungen zur Polymorphie und Kristallchemie von Silikaten der Zusammensetzung $Me_2Si_2O_5$ (Me:Na, K). 139 pages, Bremen, 2003.

**No. 208** **Pfeifer, K.**
Auswirkungen frühdiagenetischer Prozesse auf Calcit- und Barytgehalte in marinen Oberflächen-sedimenten. 110 pages, Bremen, 2003.

**No. 209** **Heuer, V.**
Spurenelemente in Sedimenten des Südatlantik. Primärer Eintrag und frühdiagenetische Überprägung. 136 pages, Bremen, 2003.

**No. 210** **Streng, M.**
Phylogenetic Aspects and Taxonomy of Calcareous Dinoflagellates. 157 pages, Bremen 2003.

**No. 211** **Boeckel, B.**
Present and past coccolith assemblages in the South Atlantic: implications for species ecology, carbonate contribution and palaeoceanographic applicability. 157 pages, Bremen, 2003.

**No. 212** **Precht, E.**
Advective interfacial exchange in permeable sediments driven by surface gravity waves and its ecological consequences. 131 pages, Bremen, 2003.

**No. 213** **Frenz, M.**
Grain-size composition of Quaternary South Atlantic sediments and its paleoceanographic significance. 123 pages, Bremen, 2003.

**No. 214** **Meggers, H. and cruise participants**
Report and preliminary results of METEOR Cruise M 53/1, Limassol - Las Palmas – Mindelo, 30.03.2002 - 03.05.2002. 81 pages, Bremen, 2003.

**No. 215** **Schulz, H.D. and cruise participants**
Report and preliminary results of METEOR Cruise M 58/1, Dakar – Las Palmas, 15.04..2003 - 12.05.2003. Bremen, 2003.

**No. 216** **Schneider, R. and cruise participants**
Report and preliminary results of METEOR Cruise M 57/1, Cape Town – Walvis Bay, 20.01. – 08.02.2003. 123 pages, Bremen, 2003.

**No. 217** **Kallmeyer, J.**
Sulfate reduction in the deep Biosphere. 157 pages, Bremen, 2003.

**No. 218** **Røy, H.**
Dynamic Structure and Function of the Diffusive Boundary Layer at the Seafloor. 149 pages, Bremen, 2003.

**No. 219** **Pätzold, J., C. Hübscher and cruise participants**
Report and preliminary results of METEOR Cruise M 52/2&3, Istanbul – Limassol – Limassol, 04.02. – 27.03.2002. Bremen, 2003.

**No. 220** **Zabel, M. and cruise participants**
Report and preliminary results of METEOR Cruise M 57/2, Walvis Bay – Walvis Bay, 11.02. – 12.03.2003. 136 pages, Bremen 2003.

**No. 221** **Salem, M.**
Geophysical investigations of submarine prolongations of alluvial fans on the western side of the Gulf of Aqaba-Red Sea. 100 pages, Bremen, 2003.

**No. 222** **Tilch, E.**
Oszillation von Wattflächen und deren fossiles Erhaltungspotential (Spiekerooger Rückseitenwatt, südliche Nordsee). 137 pages, Bremen, 2003.

**No. 223** **Frisch, U. and F. Kockel**
Der Bremen-Knoten im Strukturnetz Nordwest-Deutschlands. Stratigraphie, Paläogeographie, Strukturgeologie. 379 pages, Bremen, 2004.

**No. 224** **Kolonic, S.**
Mechanisms and biogeochemical implications of Cenomanian/Turonian black shale formation in North Africa: An integrated geochemical, millennial-scale study from the Tarfaya-LaAyoune Basin in SW Morocco. 174 pages, Bremen, 2004. Report online available only.

**No. 225** **Panteleit, B.**
Geochemische Prozesse in der Salz- Süßwasser Übergangszone. 106 pages, Bremen, 2004.

**No. 226** **Seiter, K.**
Regionalisierung und Quantifizierung benthischer Mineralisationsprozesse. 135 pages, Bremen, 2004.

**No. 227** **Bleil, U. and cruise participants**
Report and preliminary results of METEOR Cruise M 58/2, Las Palmas – Las Palmas (Canary Islands, Spain), 15.05. – 08.06.2003. 123 pages, Bremen, 2004.

**No. 228** **Kopf, A. and cruise participants**
Report and preliminary results of SONNE Cruise SO175, Miami - Bremerhaven, 12.11 - 30.12.2003. 218 pages, Bremen, 2004.

**No. 229** **Fabian, M.**
Near Surface Tilt and Pore Pressure Changes Induced by Pumping in Multi-Layered Poroelastic Half-Spaces. 121 pages, Bremen, 2004.

**No. 230** **Segl, M. , and cruise participants**
Report and preliminary results of POSEIDON cruise 304 Galway – Lisbon, 5. – 22. Oct. 2004. 27 pages, Bremen 2004

**No. 231** **Meinecke, G. and cruise participants**
Report and preliminary results of POSEIDON Cruise 296, Las Palmas – Las Palmas, 04.04 - 14.04.2003. 42 pages, Bremen 2005.

**No. 232** **Meinecke, G. and cruise participants**
Report and preliminary results of POSEIDON Cruise 310, Las Palmas – Las Palmas, 12.04 - 26.04.2004. 49 pages, Bremen 2005.

**No. 233** **Meinecke, G. and cruise participants**
Report and preliminary results of METEOR Cruise 58/3, Las Palmas - Ponta Delgada, 11.06 - 24.06.2003. 50 pages, Bremen 2005.

**No. 234** **Feseker, T.**
Numerical Studies on Groundwater Flow in Coastal Aquifers. 219 pages. Bremen 2004.

**No. 235** **Sahling, H. and cruise participants**
Report and preliminary results of R/V POSEIDON Cruise P317/4, Istanbul-Istanbul , 16 October - 4 November 2004. 92 pages, Bremen 2004.

**No. 236** **Meinecke, G. und Fahrtteilnehmer**
Report and preliminary results of POSEIDON Cruise 305, Las Palmas (Spain) - Lisbon (Portugal), October 28th – November 6th, 2004. 43 pages, Bremen 2005.

**No. 237** **Ruhland, G. and cruise participants**
Report and preliminary results of POSEIDON Cruise 319, Las Palmas (Spain) - Las Palmas (Spain), December 6th – December 17th, 2004. 50 pages, Bremen 2005.

**No. 238** **Chang, T.S.**
Dynamics of fine-grained sediments and stratigraphic evolution of a back-barrier tidal basin of the German Wadden Sea (southern North Sea). 102 pages, Bremen 2005.

**No. 239** **Lager, T.**
Predicting the source strength of recycling materials within the scope of a seepage water prognosis by means of standardized laboratory methods. 141 pages, Bremen 2005.

**No. 240** **Meinecke, G.**
DOLAN - Operationelle Datenübertragung im Ozean und Laterales Akustisches Netzwerk in der Tiefsee. Abschlußbericht. 42 pages, Bremen 2005.

**No. 241** **Guasti, E.**
Early Paleogene environmental turnover in the southern Tethys as recorded by foraminiferal and organic-walled dinoflagellate cysts assemblages. 203 pages, Bremen 2005.

**No. 242** **Riedinger, N.**
Preservation and diagenetic overprint of geochemical and geophysical signals in ocean margin sediments related to depositional dynamics. 91 pages, Bremen 2005.

**No. 243** **Ruhland, G. and cruise participants**
Report and preliminary results of POSEIDON cruise 320, Las Palmas (Spain) - Las Palmas (Spain), March 08th - March 18th, 2005. 57 pages, Bremen 2005.

**No. 244** **Inthorn, M.**
Lateral particle transport in nepheloid layers – a key factor for organic matter distribution and quality in the Benguela high-productivity area. 127 pages, Bremen, 2006.

**No. 245** **Aspetsberger, F.**
Benthic carbon turnover in continental slope and deep sea sediments: importance of organic matter quality at different time scales. 136 pages, Bremen, 2006.

**No. 246** **Hebbeln, D. and cruise participants**
Report and preliminary results of RV SONNE Cruise SO-184, PABESIA, Durban (South Africa) – Cilacap (Indonesia) – Darwin (Australia), July 08th - September 13th, 2005. 142 pages, Bremen 2006.

**No. 247** **Ratmeyer, V. and cruise participants**
Report and preliminary results of RV METEOR Cruise M61/3. Development of Carbonate Mounds on the Celtic Continental Margin, Northeast Atlantic. Cork (Ireland) – Ponta Delgada (Portugal), 04.06. – 21.06.2004. 64 pages, Bremen 2006.

**No. 248** **Wien, K.**
Element Stratigraphy and Age Models for Pelagites and Gravity Mass Flow Deposits based on Shipboard XRF Analysis. 100 pages, Bremen 2006.

**No. 249** **Krastel, S. and cruise participants**
Report and preliminary results of RV METEOR Cruise M65/2, Dakar - Las Palmas, 04.07. - 26.07.2005. 185 pages, Bremen 2006.

**No. 250** **Heil, G.M.N.**
Abrupt Climate Shifts in the Western Tropical to Subtropical Atlantic Region during the Last Glacial. 121 pages, Bremen 2006.

**No. 251** **Ruhland, G. and cruise participants**
Report and preliminary results of POSEIDON Cruise 330, Las Palmas – Las Palmas, November 21th – December 03rd, 2005. 48 pages, Bremen 2006.

**No. 252** **Mulitza , S. and cruise participants**
Report and preliminary results of METEOR Cruise M65/1, Dakar – Dakar, 11.06.- 1.07.2005. 149 pages, Bremen 2006.

**No. 253** **Kopf, A. and cruise participants**
Report and preliminary results of POSEIDON Cruise P336, Heraklion - Heraklion, 28.04. – 17.05.2006. 127 pages, Bremen, 2006.

**No. 254** **Wefer, G. and cruise participants**
Report and preliminary results of R/V METEOR Cruise M65/3, Las Palmas - Las Palmas (Spain), July 31st - August 10th, 2005. 24 pages, Bremen 2006.

**No. 255** **Hanebuth, T.J.J. and cruise participants**
Report and first results of the POSEIDON Cruise P342 GALIOMAR, Vigo – Lisboa (Portugal), August 19th – September 06th, 2006. Distribution Pattern, Residence Times and Export of Sediments on the Pleistocene/Holocene Galician Shelf (NW Iberian Peninsula). 203 pages, Bremen, 2007.

**No. 256** **Ahke, A.**
Composition of molecular organic matter pools, pigments and proteins, in Benguela upwelling and Arctic Sediments. 192 pages, Bremen 2007.

**No. 257** **Becker, V.**
Seeper - Ein Modell für die Praxis der Sickerwasserprognose. 170 pages, Bremen 2007.

**No. 258** **Ruhland, G. and cruise participants**
Report and preliminary results of Poseidon cruise 333, Las Palmas (Spain) – Las Palmas (Spain), March 1$^{st}$ – March 10$^{th}$, 2006. 32 pages, Bremen 2007.

**No. 259** **Fischer, G., G. Ruhland and cruise participants**
Report and preliminary results of Poseidon cruise 344, leg 1 and leg 2, Las Palmas (Spain) – Las Palmas (Spain), Oct. 20$^{th}$ –Nov 2$^{nd}$ & Nov. 4$^{th}$ – Nov 13$^{th}$, 2006. 46 pages, Bremen 2007.

**No. 260** **Westphal, H. and cruise participants**
Report and preliminary results of Poseidon cruise 346, MACUMA. Las Palmas (Spain) – Las Palmas (Spain), 28.12.2006 – 15.1.2007. 49 pages, Bremen 2007.